

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
3 May 2001 (03.05.2001)

PCT

(10) International Publication Number
WO 01/30768 A1

(51) International Patent Classification⁷: C07D 239/91,
A61K 31/517, A61P 37/02

(21) International Application Number: PCT/US00/29585

(22) International Filing Date: 26 October 2000 (26.10.2000)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
60/198,253 27 October 1999 (27.10.1999) US
60/213,104 21 June 2000 (21.06.2000) US

(71) Applicant (for all designated States except US): CY-
TOKINETICS, INC. [US/US]; Suite 2, 280 East Grand
Avenue, South San Francisco, CA 94080 (US).

(72) Inventors; and


(75) Inventors/Applicants (for US only): FINER, Jeffrey, T.
[US/US]; 661 Leo Drive, Foster City, CA 94404 (US).
BERGNES, Gustave [US/US]; Apartment A6, 3815 Su-
san Drive, San Bruno, CA 94066 (US). FENG, Bainian
[US/US]; 1033 Egret Street, Foster City, CA 94404 (US).
SMITH, Whitney, W. [US/US]; 1122 Richmond Street, El
Cerrito, CA 94530 (US). CHABALA, John, C. [US/US];
602 Sherwood Parkway, Mountainside, NJ 07092 (US).

(74) Agent: WEAVER, Jeffrey, K.; Beyer Weaver & Thomas,
LLP, P.O. Box 0778, Berkeley, CA 94704-0778 (US).


(81) Designated States (national): AE, AG, AL, AM, AT, AU,
AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CR, CU, CZ,
DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR,
HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR,
LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ,

[Continued on next page]

(54) Title: METHODS AND COMPOSITIONS UTILIZING QUINAZOLINONES



(57) Abstract: Quinazolinones of formulae (1a),
(1b), (1c) and (1d) are disclosed. They are useful for
treating cellular proliferative diseases and disorders
associated with KSP kinesin activity.



WO 01/30768 A1



NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM,
TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW.

(84) **Designated States (regional):** ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).

Published:

- *With international search report.*
- *Before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments.*

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

METHODS AND COMPOSITIONS UTILIZING QUINAZOLINONES

FIELD OF THE INVENTION

This invention relates to quinazolinone derivatives which are inhibitors of the mitotic kinesin KSP and are useful in the treatment of cellular proliferative diseases, for
5 example cancer, hyperplasias, restenosis, cardiac hypertrophy, immune disorders and inflammation.

BACKGROUND OF THE INVENTION

Interest in the medicinal chemistry of quinazoline derivatives was stimulated in the early 1950's with the elucidation of the structure of a quinazoline alkaloid, 3-[β -keto-
10 gamma-(3-hydroxy-2-piperidyl)-propyl]-4-quinazolone, from an Asian plant known for its antimalarial properties. In a quest to find additional antimalarial agents, various substituted quinazolines have been synthesized. Of particular import was the synthesis of the derivative 2-methyl-3-o-tolyl-4-(3H)-quinazolinone. This compound, known by the name methaqualone, though ineffective against protozoa, was found to
15 be a potent hypnotic.

Since the introduction of methaqualone and its discovery as a hypnotic, the pharmacological activity of quinazolinones and related compounds has been investigated. Quinazolinones and derivatives thereof are now known to have a wide variety of biological properties including hypnotic, sedative, analgesic, anticonvulsant,
20 antitussive and anti-inflammatory activities.

Quinazolinone derivatives for which specific biological uses have been described include U.S. Patent No. 5,147,875 describing 2-(substituted phenyl)-4-oxo-quinazolines with bronchodilator activity. U.S. Patent Nos. 3,723,432, 3,740,442, and 3,925,548 describe a class of 1-substituted-4-aryl-2(1H)-quinazolinone derivatives
25 useful as anti-inflammatory agents. European patent publication EP 0 056 637 B1 claims a class of 4(3H)-quinazolinone derivatives for the treatment of hypertension. European patent publication EP 0 884 319 A1 describes pharmaceutical compositions of quinazolin-4-one derivatives used to treat neurodegenerative, psychotropic, and drug and alcohol induced central and peripheral nervous system disorders.

Quinazolinones are among a growing number of therapeutic agents used to treat cell proliferative disorders, including cancer. For example, PCT WO 96/06616 describes a pharmaceutical composition containing a quinazolinone derivative to inhibit vascular smooth cell proliferation. PCT WO 96/19224 uses this same quinazolinone derivative to inhibit mesengial cell proliferation. U.S. Patent Nos. 4,981,856, 5,081,124 and 5,280,027 describe the use of quinazolinone derivatives to inhibit thymidylate synthase, the enzyme that catalyzes the methylation of deoxyuridine monophosphate to produce thymidine monophosphate which is required for DNA synthesis. U.S. Patent Nos. 5,747,498 and 5,773,476 describe quinazolinone derivatives used to treat cancers characterized by over-activity or inappropriate activity of tyrosine receptor kinases. U.S. Patent No. 5,037,829 claims (1H-azol-1-ylmethyl) substituted quinazoline compositions to treat carcinomas which occur in epithelial cells. PCT WO 98/34613 describes a composition containing a quinazolinone derivative useful for attenuating neovascularization and for treating malignancies. U.S. Patent 5,187,167 describes pharmaceutical compositions comprising quinazolin-4-one derivatives which possess anti-tumor activity.

Other therapeutic agents used to treat cancer include the taxanes and vinca alkaloids. Taxanes and vinca alkaloids act on microtubules, which are present in a variety of cellular structures. Microtubules are the primary structural element of the mitotic spindle. The mitotic spindle is responsible for distribution of replicate copies of the genome to each of the two daughter cells that result from cell division. It is presumed that disruption of the mitotic spindle by these drugs results in inhibition of cancer cell division, and induction of cancer cell death. However, microtubules form other types of cellular structures, including tracks for intracellular transport in nerve processes. Because these agents do not specifically target mitotic spindles, they have side effects that limit their usefulness.

Improvements in the specificity of agents used to treat cancer is of considerable interest because of the therapeutic benefits which would be realized if the side effects associated with the administration of these agents could be reduced. Traditionally, dramatic improvements in the treatment of cancer are associated with identification of therapeutic agents acting through novel mechanisms. Examples of this include not

only the taxanes, but also the camptothecin class of topoisomerase I inhibitors. From both of these perspectives, mitotic kinesins are attractive targets for new anti-cancer agents.

Mitotic kinesins are enzymes essential for assembly and function of the mitotic spindle, but are not generally part of other microtubule structures, such as in nerve processes. Mitotic kinesins play essential roles during all phases of mitosis. These enzymes are "molecular motors" that transform energy released by hydrolysis of ATP into mechanical force which drives the directional movement of cellular cargoes along microtubules. The catalytic domain sufficient for this task is a compact structure of approximately 340 amino acids. During mitosis, kinesins organize microtubules into the bipolar structure that is the mitotic spindle. Kinesins mediate movement of chromosomes along spindle microtubules, as well as structural changes in the mitotic spindle associated with specific phases of mitosis. Experimental perturbation of mitotic kinesin function causes malformation or dysfunction of the mitotic spindle, frequently resulting in cell cycle arrest and cell death.

Among the mitotic kinesins which have been identified is KSP. KSP belongs to an evolutionarily conserved kinesin subfamily of plus end-directed microtubule motors that assemble into bipolar homotetramers consisting of antiparallel homodimers. During mitosis KSP associates with microtubules of the mitotic spindle. Microinjection of antibodies directed against KSP into human cells prevents spindle pole separation during prometaphase, giving rise to monopolar spindles and causing mitotic arrest and induction of programmed cell death. KSP and related kinesins in other, non-human, organisms, bundle antiparallel microtubules and slide them relative to one another, thus forcing the two spindle poles apart. KSP may also mediate in anaphase B spindle elongation and focussing of microtubules at the spindle pole.

Human KSP (also termed HsEg5) has been described [Blangy, et al., Cell, 83:1159-69 (1995); Whitehead, et al., Arthritis Rheum., 39:1635-42 (1996); Galgio et al., J. Cell Biol., 135:339-414 (1996); Blangy, et al., J Biol. Chem., 272:19418-24 (1997); Blangy, et al., Cell Motil Cytoskeleton, 40:174-82 (1998); Whitehead and Rattner, J. Cell Sci., 111:2551-61 (1998); Kaiser, et al., JBC 274:18925-31 (1999); GenBank accession numbers: X85137, NM004523 and U37426], and a fragment of the KSP


gene (TRIP5) has been described [Lee, et al., Mol Endocrinol., 9:243-54 (1995); GenBank accession number L40372]. *Xenopus* KSP homologs (Eg5), as well as *Drosophila* KLP61 F/KRP1 30 have been reported.

Mitotic kinesins are attractive targets for the discovery and development of novel
5 mitotic chemotherapeutics. Accordingly, it is an object of the present invention to provide methods and compositions useful in the inhibition of KSP, a mitotic kinesin.

SUMMARY OF THE INVENTION

In accordance with the objects outlined above, the present invention provides compositions and methods that can be used to treat diseases of proliferating cells. The
10 compositions are KSP inhibitors, particularly human KSP inhibitors.

In one aspect, the invention relates to methods for treating cellular proliferative diseases, for treating disorders associated with KSP kinesin activity, and for inhibiting KSP kinesin. The methods employ compounds chosen from the group consisting of:



wherein:

R₁ is chosen from hydrogen, alkyl, aryl, alkylaryl, heteroaryl, alkylheteroaryl, substituted alkyl, substituted aryl, substituted alkylaryl, substituted heteroaryl, and substituted alkylheteroaryl;

5 R₂ and R₂' are independently chosen from hydrogen, alkyl, oxaalkyl, aryl, alkylaryl, heteroaryl, alkylheteroaryl, substituted alkyl, substituted aryl, substituted alkylaryl, substituted heteroaryl, and substituted alkylheteroaryl; or R₂ and R₂' taken together form a 3- to 7-membered ring;

10 R₃ is chosen from hydrogen, alkyl, aryl, alkylaryl, heteroaryl, alkylheteroaryl, substituted alkyl, substituted aryl, substituted alkylaryl, substituted heteroaryl, substituted alkylheteroaryl, oxaalkyl, oxaalkylaryl, substituted oxaalkylaryl, R₁₅O- and R₁₅-NH-;

15 R₃ is chosen from hydrogen, alkyl, aryl, alkylaryl, heteroaryl, alkylheteroaryl, substituted alkyl, substituted aryl, substituted alkylaryl, substituted heteroaryl, substituted alkylheteroaryl and R₁₅-NH-;

R₃ is chosen from alkyl, aryl, alkylaryl, heteroaryl, alkylheteroaryl, substituted alkyl, substituted aryl, substituted alkylaryl, substituted heteroaryl, and substituted alkylheteroaryl;

20 R₄ is chosen from hydrogen, alkyl, aryl, alkylaryl, heteroaryl, alkylheteroaryl, substituted alkyl, substituted aryl, substituted alkylaryl, substituted heteroaryl, substituted alkylheteroaryl, and R₁₆-alkylene-;

25 R₅, R₆, R₇ and R₈ are independently chosen from hydrogen, alkyl, alkoxy, halogen, fluoroalkyl, nitro, dialkylamino, alkylsulfonyl, alkylsulfonamido, sulfonamidoalkyl, sulfonamidoaryl, alkylthio, carboxyalkyl, carboxamido, aminocarbonyl, aryl and heteroaryl;

R₁₅ is chosen from alkyl, aryl, alkylaryl, heteroaryl, alkylheteroaryl, substituted alkyl, substituted aryl, substituted alkylaryl, substituted heteroaryl, and substituted alkylheteroaryl;

30 R₁₆ is chosen from alkoxy, amino, alkylamino, dialkylamino, N-heterocyclyl and substituted N-heterocyclyl.

Diseases and disorders that respond to therapy with compounds of the invention include cancer, hyperplasia, restenosis, cardiac hypertrophy, immune disorders and inflammation.

In another aspect, the invention relates to compounds useful in inhibiting KSP kinesin.

5 The compounds have the structures shown above.

In an additional aspect, the present invention provides methods of screening for compounds that will bind to a KSP kinesin, for example compounds that will displace or compete with the binding of the compositions of the invention. The methods comprise combining a labeled compound of the invention, a KSP kinesin, and at least
10 one candidate agent and determining the binding of the candidate bioactive agent to the KSP kinesin.

In a further aspect, the invention provides methods of screening for modulators of KSP kinesin activity. The methods comprise combining a composition of the invention, a KSP kinesin, and at least one candidate agent and determining the effect
15 of the candidate bioactive agent on the KSP kinesin activity.

BRIEF DESCRIPTION OF THE DRAWINGS

Figure 1 depicts a generic synthetic scheme to make compositions of the invention.

Figure 2 depicts a synthetic route for the synthesis of quinazolinone KSP inhibitors.

Figure 3 depicts representative chemical structures of quinazolinone KSP inhibitors.

5 Figure 4 depicts a synthetic route to substantially pure single enantiomers.

Figure 5 depicts synthetic routes to sulfonamides (5a), carbamates (5b), ureas (5c) and amines (5d).


DETAILED DESCRIPTION OF THE INVENTION

The present invention is directed to a class of novel compounds, based on a core
10 quinazolinone structure, that are modulators of mitotic kinesins. By inhibiting or
modulating mitotic kinesins, but not other kinesins (e.g., transport kinesins), specific
inhibition of cellular proliferation is accomplished. Thus, the present invention
capitalizes on the finding that perturbation of mitotic kinesin function causes
malformation or dysfunction of mitotic spindles, frequently resulting in cell cycle
15 arrest and cell death. The methods of inhibiting a human KSP kinesin comprise
contacting an inhibitor of the invention with a KSP kinesin, particularly human KSP
kinesins, including fragments and variants of KSP. The inhibition can be of the ATP
hydrolysis activity of the KSP kinesin and/or the mitotic spindle formation activity,
such that the mitotic spindles are disrupted. Meiotic spindles may also be disrupted.

20 An object of the present invention is to develop inhibitors and modulators of mitotic
kinesins, in particular KSP, for the treatment of disorders associated with cell
proliferation. Traditionally, dramatic improvements in the treatment of cancer, one
type of cell proliferative disorder, have been associated with identification of
therapeutic agents acting through novel mechanisms. Examples of this include not
25 only the taxane class of agents that appear to act on microtubule formation, but also
the camptothecin class of topoisomerase I inhibitors. The compositions and methods
described herein can differ in their selectivity and are preferably used to treat diseases

of proliferating cells, including, but not limited to cancer, hyperplasias, restenosis, cardiac hypertrophy, immune disorders and inflammation.


Accordingly, the present invention relates to methods employing quinazolinone amides of formula 1a:



5

1a


quinazolinone sulfonamides of formula 1b




1b

10

and quinazolinone amines of formulae 1c and 1d



1c



1d

wherein:

- R_1 is chosen from hydrogen, alkyl, aryl, alkylaryl, heteroaryl, alkylheteroaryl,
 5 substituted alkyl, substituted aryl, substituted alkylaryl, substituted heteroaryl,
 and substituted alkylheteroaryl;
- R_2 and R_2' are independently chosen from hydrogen, alkyl, oxaalkyl, aryl, alkylaryl,
 heteroaryl, alkylheteroaryl, substituted alkyl, substituted aryl, substituted
 alkylaryl, substituted heteroaryl, and substituted alkylheteroaryl; or R_2 and R_2'
 10 taken together form a 3- to 7-membered ring;
- R_3 is chosen from hydrogen, alkyl, aryl, alkylaryl, heteroaryl, alkylheteroaryl,
 substituted alkyl, substituted aryl, substituted alkylaryl, substituted heteroaryl,
 substituted alkylheteroaryl, oxaalkyl, oxaalkylaryl, substituted oxaalkylaryl,
 $R_{15}O-$ and $R_{15}-NH-$;
- 15 $R_{3'}$ is chosen from hydrogen, alkyl, aryl, alkylaryl, heteroaryl, alkylheteroaryl,
 substituted alkyl, substituted aryl, substituted alkylaryl, substituted heteroaryl,
 substituted alkylheteroaryl, and $R_{15}-NH-$;
- $R_{3''}$ is chosen from alkyl, aryl, alkylaryl, heteroaryl, alkylheteroaryl, substituted alkyl,
 substituted aryl, substituted alkylaryl, substituted heteroaryl, and substituted
 20 alkylheteroaryl;
- R_4 is chosen from hydrogen, alkyl, aryl, alkylaryl, heteroaryl, alkylheteroaryl,
 substituted alkyl, substituted aryl, substituted alkylaryl, substituted heteroaryl,
 substituted alkylheteroaryl, and R_{16} -alkylene-;
- R_5 , R_6 , R_7 and R_8 are independently chosen from hydrogen, alkyl, alkoxy, halogen,
 25 fluoroalkyl, nitro, dialkylamino, alkylsulfonyl, alkylsulfonamido,
 sulfonamidoalkyl, sulfonamidoaryl, alkylthio, carboxyalkyl, carboxamido,

aminocarbonyl, aryl and heteroaryl;

R₁₅ is chosen from alkyl, aryl, alkylaryl, heteroaryl, alkylheteroaryl, substituted alkyl, substituted aryl, substituted alkylaryl, substituted heteroaryl, and substituted alkylheteroaryl;

- 5 R₁₆ is chosen from alkoxy, amino, alkylamino, dialkylamino, N-heterocyclyl and substituted N-heterocyclyl.

All of the compounds falling within the foregoing parent genus and its subgenera are useful as kinesin inhibitors, but not all the compounds are novel. In particular, certain ureas (i.e. compounds in which R₃ is R₁₅NH) are disclosed in US patent 5,756,502 as
10 agents which modify cholecystokinin action. The specific exceptions in the claims reflect applicants' intent to avoid claiming subject matter that, while functionally part of the inventive concept, is not patentable to them for reasons having nothing to do with the scope of the invention.

Definitions

- 15 Alkyl is intended to include linear, branched, or cyclic hydrocarbon structures and combinations thereof. Lower alkyl refers to alkyl groups of from 1 to 5 carbon atoms. Examples of lower alkyl groups include methyl, ethyl, propyl, isopropyl, butyl, s-and t-butyl and the like. Preferred alkyl groups are those of C₂₀ or below. More preferred alkyl groups are those of C₁₃ or below. Cycloalkyl is a subset of alkyl and includes
20 cyclic hydrocarbon groups of from 3 to 13 carbon atoms. Examples of cycloalkyl groups include c-propyl, c-butyl, c-pentyl, norbornyl, adamantyl and the like. In this application, alkyl refers to alkanyl, alkenyl and alkynyl residues; it is intended to include cyclohexylmethyl, vinyl, allyl, isoprenyl and the like. Alkylene refers to the same residues as alkyl, but having two points of attachment. Examples of alkylene
25 include ethylene (-CH₂CH₂-), propylene (-CH₂CH₂CH₂-), dimethylpropylene (-CH₂C(CH₃)₂CH₂-) and cyclohexylpropylene (-CH₂CH₂CH(C₆H₁₃)-). When an alkyl residue having a specific number of carbons is named, all geometric isomers having that number of carbons are intended to be encompassed; thus, for example, "butyl" is meant to include n-butyl, sec-butyl, isobutyl and t-butyl; "propyl" includes n-propyl
30 and isopropyl.

Alkoxy or alkoxyl refers to groups of from 1 to 8 carbon atoms of a straight, branched, cyclic configuration and combinations thereof attached to the parent structure through an oxygen. Examples include methoxy, ethoxy, propoxy, isopropoxy, cyclopropyloxy, cyclohexyloxy and the like. Lower-alkoxy refers to groups
5 containing one to four carbons.

Acyl refers to groups of from 1 to 8 carbon atoms of a straight, branched, cyclic configuration, saturated, unsaturated and aromatic and combinations thereof, attached to the parent structure through a carbonyl functionality. One or more carbons in the acyl residue may be replaced by nitrogen, oxygen or sulfur as long as the point of
10 attachment to the parent remains at the carbonyl. Examples include acetyl, benzoyl, propionyl, isobutyryl, t-butoxycarbonyl, benzyloxycarbonyl and the like. Lower-acyl refers to groups containing one to four carbons.

Aryl and heteroaryl mean a 5- or 6-membered aromatic or heteroaromatic ring containing 0-3 heteroatoms selected from O, N, or S; a bicyclic 9- or 10-membered
15 aromatic or heteroaromatic ring system containing 0-3 heteroatoms selected from O, N, or S; or a tricyclic 13- or 14-membered aromatic or heteroaromatic ring system containing 0-3 heteroatoms selected from O, N, or S. The aromatic 6- to 14-membered carbocyclic rings include, e.g., benzene, naphthalene, indane, tetralin, and fluorene and the 5- to 10-membered aromatic heterocyclic rings include, e.g.,
20 imidazole, pyridine, indole, thiophene, benzopyranone, thiazole, furan, benzimidazole, quinoline, isoquinoline, quinoxaline, pyrimidine, pyrazine, tetrazole and pyrazole.

Alkylaryl refers to a residue in which an aryl moiety is attached to the parent structure via an alkyl residue. Examples are benzyl, phenethyl, phenylvinyl, phenylallyl and the like. Oxaalkyl and oxaalkylaryl refer to alkyl and alkylaryl residues in which one or
25 more methylenes have been replaced by oxygen. Examples of oxaalkyl and oxaalkylaryl residues are ethoxyethoxyethyl (3,6-dioxaoctyl), benzyloxymethyl and phenoxymethyl; in general, glycol ethers, such as polyethyleneglycol, are intended to be encompassed by this group. Alkylheteroaryl refers to a residue in which a heteroaryl moiety is attached to the parent structure via an alkyl residue. Examples
30 include furanylmethyl, pyridinylmethyl, pyrimidinylethyl and the like.

Heterocycle means a cycloalkyl or aryl residue in which one to four of the carbons is replaced by a heteroatom such as oxygen, nitrogen or sulfur. Examples of heterocycles that fall within the scope of the invention include imidazoline, pyrrolidine, pyrazole, pyrrole, indole, quinoline, isoquinoline, tetrahydroisoquinoline, benzofuran, benzodioxan, benzodioxole (commonly referred to as methylenedioxyphenyl, when occurring as a substituent), tetrazole, morpholine, thiazole, pyridine, pyridazine, pyrimidine, thiophene, furan, oxazole, oxazoline, isoxazole, dioxane, tetrahydrofuran and the like. "N-heterocyclyl" refers to a nitrogen-containing heterocycle as a substituent residue. The term heterocyclyl encompasses heteroaryl, which is a subset of heterocyclyl. Examples of N-heterocyclyl residues include 4-morpholinyl, 4-thiomorpholinyl, 1-piperidinyl, 1-pyrrolidinyl, 3-thiazolidinyl, piperazinyl and 4-(3,4-dihydrobenzoxazinyl). Examples of substituted heterocyclyl include 4-methyl-1-piperazinyl and 4-benzyl-1-piperidinyl.

Substituted alkyl, aryl and heteroaryl refer to alkyl, aryl or heteroaryl wherein H atoms are replaced with alkyl, halogen, hydroxy, alkoxy, alkylendioxy (e.g. methylenedioxy) fluoroalkyl, carboxy (-COOH), carboalkoxy (i.e. acyloxy RCOO-), carboxyalkyl (-COOR), carboxamido, sulfonamidoalkyl, sulfonamidoaryl, aminocarbonyl, benzyloxycarbonylamino (CBZ-amino), cyano, carbonyl, nitro, dialkylamino, alkylamino, amino, alkylthio, alkylsulfinyl, alkylsulfonyl, alkylsulfonamido, arylthio, arylsulfinyl, arylsulfonyl, amidino, phenyl, benzyl, heteroaryl, heterocyclyl, phenoxy, benzyloxy, or heteroaryloxy. For the purposes of the present invention, substituted alkyl also includes oxaalkyl residues, i.e. alkyl residues in which one or more carbons has been replaced by oxygen.

Halogen refers to fluorine, chlorine, bromine or iodine. Fluorine, chlorine and bromine are preferred. Dihaloaryl, dihaloalkyl, trihaloaryl etc. refer to aryl and alkyl substituted with a plurality of halogens, but not necessarily a plurality of the same halogen; thus 4-chloro-3-fluorophenyl is within the scope of dihaloaryl.

Most of the compounds described herein contain one or more asymmetric centers (e.g. the carbon to which R₂ and R₂' are attached) and may thus give rise to enantiomers, diastereomers, and other stereoisomeric forms that may be defined, in terms of absolute stereochemistry, as (R)- or (S)-. The present invention is meant to include all

such possible isomers, including racemic mixtures, optically pure forms and intermediate mixtures. Optically active (R)- and (S)- isomers may be prepared using chiral synthons or chiral reagents, or resolved using conventional techniques. When the compounds described herein contain olefinic double bonds or other centers of geometric asymmetry, and unless specified otherwise, it is intended that the compounds include both E and Z geometric isomers. Likewise, all tautomeric forms are also intended to be included.

When desired, the R- and S-isomers may be resolved by methods known to those skilled in the art, for example by formation of diastereoisomeric salts or complexes which may be separated, for example, by crystallisation; via formation of diastereoisomeric derivatives which may be separated, for example, by crystallisation, gas-liquid or liquid chromatography; selective reaction of one enantiomer with an enantiomer-specific reagent, for example enzymatic oxidation or reduction, followed by separation of the modified and unmodified enantiomers; or gas-liquid or liquid chromatography in a chiral environment, for example on a chiral support, such as silica with a bound chiral ligand or in the presence of a chiral solvent. It will be appreciated that where the desired enantiomer is converted into another chemical entity by one of the separation procedures described above, a further step may be required to liberate the desired enantiomeric form. Alternatively, specific enantiomer may be synthesized by asymmetric synthesis using optically active reagents, substrates, catalysts or solvents, or by converting one enantiomer to the other by asymmetric transformation. An example of a synthesis from optically active starting materials is shown in Figure 4.

In one embodiment, as will be appreciated by those in the art, the two adjacent R_2 groups may be fused together to form a ring structure. Again, the fused ring structure may contain heteroatoms and may be substituted with one or more substitution groups "R". It should additionally be noted that for cycloalkyl (i.e. saturated ring structures), each position may contain two substitution groups, R and R'.

Considering structures 1a, 1b, 1c and 1d, but focusing on 1a, in a preferred embodiment R_1 is selected from hydrogen, alkyl, aryl, substituted alkyl, substituted aryl, heteroaryl, substituted heteroaryl, alkylaryl and substituted alkylaryl.

In a more preferred embodiment R_1 is selected from hydrogen, lower alkyl, substituted lower alkyl, benzyl, substituted benzyl, phenyl, naphthyl and substituted phenyl.

In a most preferred embodiment R_1 is chosen from hydrogen, ethyl, propyl, methoxyethyl, naphthyl, phenyl, bromophenyl, chlorophenyl, methoxyphenyl, ethoxyphenyl, tolyl, dimethylphenyl, chlorofluorophenyl, methylchlorophenyl, ethylphenyl, phenethyl, benzyl, chlorobenzyl, methylbenzyl, methoxybenzyl, cyanobenzyl, hydroxybenzyl, tetrahydrofuranylmethyl and (ethoxycarbonyl)ethyl.

In a preferred embodiment R_2 is hydrogen, alkyl or substituted alkyl. As will be appreciated by those in the art, Structures 1a, 1b, 1c and 1d possess a potentially chiral center at the carbon to which R_2 is attached. Thus, the R_2 position may comprise two substitution groups, R_2 and R_2' . The R_2 and R_2' groups may be the same or different; if different, the composition is chiral. When the R_2 and R_2' are different, preferred embodiments utilize only a single non-hydrogen R_2 . The invention contemplates the use of pure enantiomers and mixtures of enantiomers, including racemic mixtures, although the use of the substantially optically pure enantiomer will generally be preferred.

In a more preferred embodiment, R_2 is chosen from hydrogen, lower alkyl and substituted lower alkyl, and R_2' is hydrogen. In a most preferred embodiment R_2 is chosen from hydrogen, methyl, ethyl, propyl, methylthioethyl, aminobutyl, (CBZ)aminobutyl, cyclohexylmethyl, benzyloxymethyl, methylsulfinylethyl, methylsulfinylmethyl, hydroxymethyl, benzyl and indolylmethyl.

In a preferred embodiment R_3 is selected from alkyl, substituted alkyl, alkylaryl, heteroaryl, aryl, substituted aryl, substituted oxaalkylaryl, $R_{15}O-$ and $R_{15}NH-$, and R_{15} is chosen from alkyl, aryl and substituted aryl.

In a more preferred embodiment, when R_3 is not $R_{15}NH$, R_3 is chosen from C_1-C_{13} alkyl; substituted lower alkyl; phenyl; naphthyl; phenyl substituted with one or more halo, lower alkyl, loweralkoxy, nitro, carboxy, methylenedioxy or trifluoromethyl; biphenyl; benzyl; phenoxymethyl; halophenoxymethyl; phenylvinyl; heteroaryl; heteroaryl substituted with lower alkyl; and benzyloxymethyl.

In a most preferred embodiment, when R_3 is not $R_{15}NH$, R_3 is chosen from ethyl, propyl, chloropropyl, butoxy, heptyl, butyl, octyl, tridecanyl, (ethoxycarbonyl)ethyl, dimethylaminoethyl, dimethylaminomethyl, phenyl, naphthyl, halophenyl, dihalophenyl, cyanophenyl, halo(trifluoromethyl)phenyl, chlorophenoxymethyl, methoxyphenyl, carboxyphenyl, ethylphenyl, tolyl, biphenyl, methylenedioxyphenyl, methylsulfonylphenyl, methoxychlorophenyl, chloronaphthyl, methylhalophenyl, trifluoromethylphenyl, butylphenyl, pentylphenyl, methylnitrophenyl, phenoxymethyl, dimethoxyphenyl, phenylvinyl, nitrochlorophenyl, nitrophenyl, dinitrophenyl, bis(trifluoromethyl)phenyl, benzyloxymethyl, benzyl, furanyl, benzofuranyl, pyridinyl, indolyl, methylpyridinyl, quinolinyl, picolinyl, pyrazolyl, and imidazolyl.

In a more preferred embodiment, when R_3 is $R_{15}NH$, R_{15} is chosen from lower alkyl; cyclohexyl; phenyl; and phenyl substituted with halo, lower alkyl, loweralkoxy, or lower alkylthio.

In a most preferred embodiment, when R_3 is $R_{15}NH$, R_{15} is isopropyl, butyl, cyclohexyl, phenyl, bromophenyl, dichlorophenyl, methoxyphenyl, ethylphenyl, tolyl, trifluoromethylphenyl or methylthiophenyl.

In a preferred embodiment R_4 is chosen from alkyl, aryl, alkylaryl, alkylheteroaryl, substituted alkyl, substituted aryl, and R_{16} -alkylene-, and R_{16} is chosen from alkoxy, amino, alkylamino, dialkylamino and N-heterocyclyl.

In a more preferred embodiment, R_4 is selected from lower alkyl, substituted lower alkyl, cyclohexyl; phenyl substituted with hydroxy, lower alkoxy or lower alkyl; benzyl; heteroarylmethyl; heteroarylethyl; heteroarylpropyl and R_{16} -alkylene-, wherein R_{16} is amino, lower alkylamino, di(lower alkyl)amino, lower alkoxy, or N-heterocyclyl.

In a most preferred embodiment, R_4 is chosen from methyl, ethyl, propyl, butyl, cyclohexyl, carboxyethyl, carboxymethyl, methoxyethyl, hydroxyethyl, hydroxypropyl, dimethylaminoethyl, dimethylaminopropyl, diethylaminoethyl, diethylaminopropyl, aminopropyl, methylaminopropyl, 2,2-dimethyl-3-(dimethylamino)propyl, 1-cyclohexyl-4-(diethylamino)butyl, aminoethyl, aminobutyl, aminopentyl, aminohexyl, aminoethoxyethyl, isopropylaminopropyl,

diisopropylaminoethyl, 1-methyl-4-(diethylamino)butyl, (t-Boc)aminopropyl,
 hydroxyphenyl, benzyl, methoxyphenyl, methylmethoxyphenyl, dimethylphenyl, tolyl,
 ethylphenyl, (oxopyrrolidinyl)propyl, (methoxycarbonyl)ethyl, benzylpiperidinyl,
 pyridinylethyl, pyridinylmethyl, morpholinylethyl morpholinylpropyl, piperidinyl,
 5 azetidinylmethyl, azetidinypropyl pyrrolidinylethyl, pyrrolidinylpropyl,
 piperidinylmethyl, piperidinylethyl, imidazolylpropyl, imidazolylethyl,
 (ethylpyrrolidinyl)methyl, (methylpyrrolidinyl)ethyl, (methylpiperidinyl)propyl,
 (methylpiperazinyl)propyl, furanylmethyl and indolylethyl.

In other preferred embodiments R_5 is hydrogen or halo; R_6 is hydrogen, methyl or
 10 halo; R_7 is hydrogen, halo, methyl or trifluoromethyl; and R_8 is hydrogen or halo.

In a particularly preferred subgenus, R_1 is benzyl or halobenzyl; R_2 is chosen from
 ethyl and propyl; R_2' is hydrogen; R_3 is substituted phenyl; $R_{3'}$ is substituted phenyl;
 $R_{3''}$ is substituted phenyl; R_4 is $-(CH)_mOH$ or $-(CH_2)_pR_{16}$; m is two or three; p is one
 to three; R_5 is hydrogen; R_6 is hydrogen; R_7 is halo; R_8 is hydrogen; and R_{16} is chosen
 15 from amino, propylamino, and azetidinyl.

When considering primarily the sulfonamides of structure 1b, R_1 is preferably chosen
 from hydrogen, lower alkyl, substituted lower alkyl, benzyl, substituted benzyl, phenyl
 and substituted phenyl; R_2 is chosen from hydrogen and lower alkyl and R_2' is
 hydrogen; $R_{3'}$ is chosen from C_1 - C_{13} alkyl; phenyl; naphthyl; phenyl substituted with
 20 halo, lower alkyl, lower alkoxy, nitro, methylenedioxy, or trifluoromethyl; biphenyl
 and heteroaryl; and R_4 is chosen from lower alkyl, cyclohexyl; phenyl substituted with
 hydroxy, lower alkoxy or lower alkyl; benzyl; heteroarylmethyl; heteroarylethyl;
 heteroarylpropyl; heteroarylethyl; heteroarylpropyl and R_{16} -alkylene, wherein R_{16} is
 di(lower alkyl)amino, (lower alkyl)amino, amino, lower alkoxy, or N-heterocyclyl,
 25 particularly pyrrolidino, piperidino or imidazolyl.

When considering primarily the sulfonamides of structure 1b, R_1 is most preferably
 chosen from lower alkyl, benzyl, substituted benzyl and substituted phenyl; R_2 is
 hydrogen or lower alkyl; R_2' is hydrogen; R_3 is chosen from substituted phenyl and
 naphthyl; R_4 is R_{16} -alkylene-; R_7 is hydrogen, fluoro, methyl or chloro; R_5 , R_6 and R_8
 30 are hydrogen; and R_{16} is chosen from di(lower alkylamino), (lower alkyl)amino,

amino, pyrrolidino, piperidino, imidazolyl and morpholino.

When considering primarily the amines of structures 1c and 1d, R₁ is preferably chosen from hydrogen, lower alkyl, substituted lower alkyl, benzyl, substituted benzyl, phenyl, naphthyl and substituted phenyl; R₂ is chosen from hydrogen, lower alkyl and substituted lower alkyl and R₂' is hydrogen; R₃ is chosen from C₁-C₁₃ alkyl; substituted lower alkyl; phenyl; naphthyl; phenyl substituted with halo, lower alkyl, lower alkoxy, nitro, methylenedioxy, or trifluoromethyl; biphenyl, benzyl and heterocyclyl; and R₄ is chosen from lower alkyl; cyclohexyl; phenyl substituted with hydroxy, lower alkoxy or lower alkyl; benzyl; substituted benzyl; heterocyclyl; heteroarylmethyl; heteroarylethyl; heteroarylpropyl and R₁₆-alkylene, wherein R₁₆ is di(lower alkyl)amino, (lower alkyl)amino, amino, lower alkoxy, or N-heterocyclyl.

When considering primarily the amines of structure 1c and 1d, R₁ is most preferably chosen from lower alkyl, benzyl, substituted benzyl and substituted phenyl; R₂ is hydrogen or lower alkyl; R₂' is hydrogen; R₃ is chosen from substituted phenyl, heterocyclyl and naphthyl; R₄ is chosen from substituted benzyl, heterocyclyl and R₁₆-alkylene-; R₆ and R₇ are chosen from hydrogen and halo; R₅ and R₈ are hydrogen; and R₁₆ is chosen from di(lower alkylamino), (lower alkyl)amino, amino, pyrrolidinyl, piperidinyl, imidazolyl and morpholinyl. When R₃ is present (as in 1d) it is most preferably chosen from halophenyl, polyhalophenyl, tolyl, dimethylphenyl, methoxyphenyl, dimethoxyphenyl, cyanophenyl, trifluoromethylphenyl, trifluoromethoxyphenyl, bis(trifluoromethyl)phenyl, carboxyphenyl, t-butylphenyl, methoxycarbonylphenyl, piperidinyl and naphthyl.

The compositions of the invention are synthesized as outlined below, utilizing techniques well known in the art. For example, as described in Ager et al., J. of Med. Chem.,

20:379-386 (1977), hereby incorporated by reference, quinazolinones can be obtained by acid-catalyzed condensation of N-acylanthranilic acids with aromatic primary amines. Other processes for preparing quinazolinones are described in U.S. Patent applications 5,783,577, 5,922,866 and 5,187,167, all of which are incorporated by reference.

The compositions of the invention may be made as shown in Figures 1, 2, 4 and 5. Compounds of formulae 1d are made in analogous fashion to figure 1, except that the acyl halide in the final step is replaced by an alkyl halide.

Once made, the compositions of the invention find use in a variety of applications. As
5 will be appreciated by those in the art, mitosis may be altered in a variety of ways; that is, one can affect mitosis either by increasing or decreasing the activity of a component in the mitotic pathway. Stated differently, mitosis may be affected (e.g., disrupted) by disturbing equilibrium, either by inhibiting or activating certain components. Similar approaches may be used to alter meiosis.

10 In a preferred embodiment, the compositions of the invention are used to modulate mitotic spindle formation, thus causing prolonged cell cycle arrest in mitosis. By "modulate" herein is meant altering mitotic spindle formation, including increasing and decreasing spindle formation. By "mitotic spindle formation" herein is meant organization of microtubules into bipolar structures by mitotic kinesins. By "mitotic
15 spindle dysfunction" herein is meant mitotic arrest and monopolar spindle formation.

The compositions of the invention are useful to bind to and/or modulate the activity of a mitotic kinesin, KSP. In a preferred embodiment, the KSP is human KSP, although KSP kinesins from other organisms may also be used. In this context, modulate means either increasing or decreasing spindle pole separation, causing malformation,
20 i.e., splaying, of mitotic spindle poles, or otherwise causing morphological perturbation of the mitotic spindle. Also included within the definition of KSP for these purposes are variants and/or fragments of KSP. See U.S. Patent Application "Methods of Screening for Modulators of Cell Proliferation and Methods of Diagnosing Cell Proliferation States", filed Oct. 27, 1999 (U.S. Serial Number
25 09/428,156), hereby incorporated by reference in its entirety. In addition, other mitotic kinesins may be used in the present invention. However, the compositions of the invention have been shown to have specificity for KSP.

For assay of activity, generally either KSP or a compound according to the invention is non-diffusably bound to an insoluble support having isolated sample receiving areas
30 (e.g., a microtiter plate, an array, etc.). The insoluble support may be made of any

composition to which the compositions can be bound, is readily separated from soluble material, and is otherwise compatible with the overall method of screening. The surface of such supports may be solid or porous and of any convenient shape. Examples of suitable insoluble supports include microtiter plates, arrays, membranes and beads. These are typically made of glass, plastic (e.g., polystyrene), polysaccharides, nylon or nitrocellulose, Teflon™, etc. Microtiter plates and arrays are especially convenient because a large number of assays can be carried out simultaneously, using small amounts of reagents and samples. The particular manner of binding of the composition is not crucial so long as it is compatible with the reagents and overall methods of the invention, maintains the activity of the composition and is nondiffusible. Preferred methods of binding include the use of antibodies (which do not sterically block either the ligand binding site or activation sequence when the protein is bound to the support), direct binding to "sticky" or ionic supports, chemical crosslinking, the synthesis of the protein or agent on the surface, etc. Following binding of the protein or agent, excess unbound material is removed by washing. The sample receiving areas may then be blocked through incubation with bovine serum albumin (BSA), casein or other innocuous protein or other moiety.

The antimitotic agents of the invention may be used on their own to modulate the activity of a mitotic kinesin, particularly KSP. In this embodiment, the mitotic agents of the invention are combined with KSP and the activity of KSP is assayed. Kinesin activity is known in the art and includes one or more kinesin activities. Kinesin activities include the ability to affect ATP hydrolysis; microtubule binding; gliding and polymerization/depolymerization (effects on microtubule dynamics); binding to other proteins of the spindle; binding to proteins involved in cell-cycle control; serving as a substrate to other enzymes; such as kinases or proteases; and specific kinesin cellular activities such as spindle pole separation.

Methods of performing motility assays are well known to those of skill in the art. [See e.g., Hall, et al. (1996), Biophys. J., 71: 3467-3476, Turner et al., 1996, Anal Biochem. 242 (1):20-5; Gittes et al., 1996, Biophys. J. 70(1): 418-29; Shirakawa et al., 1995, J. Exp. Biol. 198: 1809-15; Winkelmann et al., 1995, Biophys. J. 68: 2444-53; Winkelmann et al., 1995, Biophys. J. 68: 72S.]

Methods known in the art for determining ATPase hydrolysis activity also can be used. Preferably, solution based assays are utilized. U.S. application 09/314,464, filed May 18, 1999, hereby incorporated by reference in its entirety, describes such assays. Alternatively, conventional methods are used. For example, P_i release from
5 kinesin can be quantified. In one preferred embodiment, the ATPase hydrolysis activity assay utilizes 0.3 M PCA (perchloric acid) and malachite green reagent (8.27 mM sodium molybdate II, 0.33 mM malachite green oxalate, and 0.8 mM Triton X-100). To perform the assay, 10 μ L of reaction is quenched in 90 μ L of cold 0.3 M PCA. Phosphate standards are used so data can be converted to mM inorganic
10 phosphate released. When all reactions and standards have been quenched in PCA, 100 μ L of malachite green reagent is added to the relevant wells in e.g., a microtiter plate. The mixture is developed for 10-15 minutes and the plate is read at an absorbance of 650 nm. If phosphate standards were used, absorbance readings can be converted to mM P_i and plotted over time. Additionally, ATPase assays known in the
15 art include the luciferase assay.

ATPase activity of kinesin motor domains also can be used to monitor the effects of modulating agents. In one embodiment ATPase assays of kinesin are performed in the absence of microtubules. In another embodiment, the ATPase assays are performed in the presence of microtubules. Different types of modulating agents can be detected in
20 the above assays. In a preferred embodiment, the effect of a modulating agent is independent of the concentration of microtubules and ATP. In another embodiment, the effect of the agents on kinesin ATPase can be decreased by increasing the concentrations of ATP, microtubules or both. In yet another embodiment, the effect of the modulating agent is increased by increasing concentrations of ATP,
25 microtubules or both.

Agents that modulate the biochemical activity of KSP in vitro may then be screened in vivo. Methods for such agents in vivo include assays of cell cycle distribution, cell viability, or the presence, morphology, activity, distribution, or amount of mitotic
30 spindles. Methods for monitoring cell cycle distribution of a cell population, for example, by flow cytometry, are well known to those skilled in the art, as are methods for determining cell viability. See for example, U.S. Patent Application "Methods of

Screening for Modulators of Cell Proliferation and Methods of Diagnosing Cell Proliferation States," filed Oct. 22, 1999, serial number 09/428,156, hereby incorporated by reference in its entirety.

5 In addition to the assays described above, microscopic methods for monitoring spindle formation and malformation are well known to those of skill in the art (see, e.g., Whitehead and Rattner (1998), J. Cell Sci. 111:2551-61; Galgio et al, (1996) J. Cell biol., 135:399-414).

The compositions of the invention inhibit the KSP kinesin. One measure of inhibition is IC_{50} , defined as the concentration of the composition at which the activity of KSP is
10 decreased by fifty percent. Preferred compositions have IC_{50} 's of less than about 1 mM, with preferred embodiments having IC_{50} 's of less than about 100 μ M, with more preferred embodiments having IC_{50} 's of less than about 10 μ M, with particularly preferred embodiments having IC_{50} 's of less than about 1 μ M, and especially preferred
15 embodiments having IC_{50} 's of less than about 100 nM, and with the most preferred embodiments having IC_{50} 's of less than about 10 nM. Measurement of IC_{50} is done using an ATPase assay.

Another measure of inhibition is K_i . For compounds with IC_{50} 's less than 1 μ M, the K_i or K_d is defined as the dissociation rate constant for the interaction of the quinazolinone with KSP. Preferred compounds have K_i 's of less than about 100 μ M,
20 with preferred embodiments having K_i 's of less than about 10 μ M, and particularly preferred embodiments having K_i 's of less than about 1 μ M and especially preferred embodiments having K_i 's of less than about 100 nM, and with the most preferred
25 embodiments having K_i 's of less than about 10 nM. The K_i for a compound is determined from the IC_{50} based on three assumptions. First, only one compound molecule binds to the enzyme and there is no cooperativity. Second, the concentrations of active enzyme and the compound tested are known (i.e., there are no significant amounts of impurities or inactive forms in the preparations). Third, the enzymatic rate of the enzyme-inhibitor complex is zero. The rate (i.e., compound concentration) data are fitted to the equation:

$$V = V_{\max} E_0 \left[I - \frac{(E_0 + I_0 + K_d) - \sqrt{(E_0 + I_0 + K_d)^2 - 4 E_0 I_0}}{2 E_0} \right]$$

Where V is the observed rate, V_{\max} is the rate of the free enzyme, I_0 is the inhibitor concentration, E_0 is the enzyme concentration, and K_d is the dissociation constant of the enzyme-inhibitor complex.

- 5 Another measure of inhibition is GI_{50} , defined as the concentration of the compound that results in a decrease in the rate of cell growth by fifty percent. Preferred compounds have GI_{50} 's of less than about 1 mM. The level of preferability of
embodiments is a function of their GI_{50} : those having GI_{50} 's of less than about 20 μ M
are more preferred; those having GI_{50} 's of 10 μ M more so; those having GI_{50} of less
10 than about 1 μ M more so; those having GI_{50} 's of 100 nM more so; those having GI_{50}
of less than about 10 nM even more so. Measurement of GI_{50} is done using a cell
proliferation assay.

The compositions of the invention are used to treat cellular proliferation diseases. Disease states which can be treated by the methods and compositions provided herein
15 include, but are not limited to, cancer (further discussed below), autoimmune disease, arthritis, graft rejection, inflammatory bowel disease, proliferation induced after medical procedures, including, but not limited to, surgery, angioplasty, and the like. It is appreciated that in some cases the cells may not be in a hyper or hypo proliferation state (abnormal state) and still require treatment. For example, during wound healing,
20 the cells may be proliferating "normally", but proliferation enhancement may be desired. Similarly, as discussed above, in the agriculture arena, cells may be in a "normal" state, but proliferation modulation may be desired to enhance a crop by directly enhancing growth of a crop, or by inhibiting the growth of a plant or organism which adversely affects the crop. Thus, in one embodiment, the invention herein
25 includes application to cells or individuals afflicted or impending affliction with any one of these disorders or states.

The compositions and methods provided herein are particularly deemed useful for the treatment of cancer including solid tumors such as skin, breast, brain, cervical carcinomas, testicular carcinomas, etc. More particularly, cancers that may be treated by the compositions and methods of the invention include, but are not limited to:

- 5 Cardiac: sarcoma (angiosarcoma, fibrosarcoma, rhabdomyosarcoma, liposarcoma), myxoma, rhabdomyoma, fibroma, lipoma and teratoma; Lung: bronchogenic carcinoma (squamous cell, undifferentiated small cell, undifferentiated large cell, adenocarcinoma), alveolar (bronchiolar) carcinoma, bronchial adenoma, sarcoma, lymphoma, chondromatous hamartoma, mesothelioma; Gastrointestinal: esophagus
- 10 (squamous cell carcinoma, adenocarcinoma, leiomyosarcoma, lymphoma), stomach (carcinoma, lymphoma, leiomyosarcoma), pancreas (ductal adenocarcinoma, insulinoma, glucagonoma, gastrinoma, carcinoid tumors, vipoma), small bowel (adenocarcinoma, lymphoma, carcinoid tumors, Kaposi's sarcoma, leiomyoma, hemangioma, lipoma, neurofibroma, fibroma), large bowel (adenocarcinoma, tubular
- 15 adenoma, villous adenoma, hamartoma, leiomyoma); Genitourinary tract: kidney (adenocarcinoma, Wilm's tumor [nephroblastoma], lymphoma, leukemia), bladder and urethra (squamous cell carcinoma, transitional cell carcinoma, adenocarcinoma), prostate (adenocarcinoma, sarcoma), testis (seminoma, teratoma, embryonal carcinoma, teratocarcinoma, choriocarcinoma, sarcoma, interstitial cell carcinoma,
- 20 fibroma, fibroadenoma, adenomatoid tumors, lipoma); Liver: hepatoma (hepatocellular carcinoma), cholangiocarcinoma, hepatoblastoma, angiosarcoma, hepatocellular adenoma, hemangioma; Bone: osteogenic sarcoma (osteosarcoma), fibrosarcoma, malignant fibrous histiocytoma, chondrosarcoma, Ewing's sarcoma, malignant lymphoma (reticulum cell sarcoma), multiple myeloma, malignant giant
- 25 cell tumor chordoma, osteochondroma (osteochondrogenous exostoses), benign chondroma, chondroblastoma, chondromyxofibroma, osteoid osteoma and giant cell tumors; Nervous system: skull (osteoma, hemangioma, granuloma, xanthoma, osteitis deformans), meninges (meningioma, meningiosarcoma, gliomatosis), brain (astrocytoma, medulloblastoma, glioma, ependymoma, germinoma [pinealoma],
- 30 glioblastoma multiform, oligodendroglioma, schwannoma, retinoblastoma, congenital tumors), spinal cord neurofibroma, meningioma, glioma, sarcoma); Gynecological: uterus (endometrial carcinoma), cervix (cervical carcinoma, pre-tumor cervical

dysplasia), ovaries (ovarian carcinoma [serous cystadenocarcinoma, mucinous cystadenocarcinoma, unclassified carcinoma], granulosa-thecal cell tumors, Sertoli-Leydig cell tumors, dysgerminoma, malignant teratoma), vulva (squamous cell carcinoma, intraepithelial carcinoma, adenocarcinoma, fibrosarcoma, melanoma),
5 vagina (clear cell carcinoma, squamous cell carcinoma, botryoid sarcoma (embryonal rhabdomyosarcoma), fallopian tubes (carcinoma); Hematologic: blood (myeloid leukemia [acute and chronic], acute lymphoblastic leukemia, chronic lymphocytic leukemia, myeloproliferative diseases, multiple myeloma, myelodysplastic syndrome), Hodgkin's disease, non-Hodgkin's lymphoma [malignant lymphoma]; Skin: malignant
10 melanoma, basal cell carcinoma, squamous cell carcinoma, Kaposi's sarcoma, moles dysplastic nevi, lipoma, angioma, dermatofibroma, keloids, psoriasis; and Adrenal glands: neuroblastoma. Thus, the term "cancerous cell" as provided herein, includes a cell afflicted by any one of the above identified conditions.

Accordingly, the compositions of the invention are administered to cells. By
15 "administered" herein is meant administration of a therapeutically effective dose of the mitotic agents of the invention to a cell either in cell culture or in a patient. By "therapeutically effective dose" herein is meant a dose that produces the effects for which it is administered. The exact dose will depend on the purpose of the treatment, and will be ascertainable by one skilled in the art using known techniques. As is
20 known in the art, adjustments for systemic versus localized delivery, age, body weight, general health, sex, diet, time of administration, drug interaction and the severity of the condition may be necessary, and will be ascertainable with routine experimentation by those skilled in the art. By "cells" herein is meant almost any cell in which mitosis or meiosis can be altered.

25 A "patient" for the purposes of the present invention includes both humans and other animals, particularly mammals, and other organisms. Thus the methods are applicable to both human therapy and veterinary applications. In the preferred embodiment the patient is a mammal, and in the most preferred embodiment the patient is human.

Mitotic agents having the desired pharmacological activity may be administered in a
30 physiologically acceptable carrier to a patient, as described herein. Depending upon the manner of introduction, the compounds may be formulated in a variety of ways as

discussed below. The concentration of therapeutically active compound in the formulation may vary from about 0.1-100 wt.%. The agents may be administered alone or in combination with other treatments, i.e., radiation, or other chemotherapeutic agents.

- 5 In a preferred embodiment, the pharmaceutical compositions are in a water soluble form, such as pharmaceutically acceptable salts, which is meant to include both acid and base addition salts. "Pharmaceutically acceptable acid addition salt" refers to those salts that retain the biological effectiveness of the free bases and that are not biologically or otherwise undesirable, formed with inorganic acids such as
- 10 hydrochloric acid, hydrobromic acid, sulfuric acid, nitric acid, phosphoric acid and the like, and organic acids such as acetic acid, propionic acid, glycolic acid, pyruvic acid, oxalic acid, maleic acid, malonic acid, succinic acid, fumaric acid, tartaric acid, citric acid, benzoic acid, cinnamic acid, mandelic acid, methanesulfonic acid, ethanesulfonic acid, p-toluenesulfonic acid, salicylic acid and the like.
- 15 "Pharmaceutically acceptable base addition salts" include those derived from inorganic bases such as sodium, potassium, lithium, ammonium, calcium, magnesium, iron, zinc, copper, manganese, aluminum salts and the like. Particularly preferred are the ammonium, potassium, sodium, calcium, and magnesium salts. Salts derived from pharmaceutically acceptable organic non-toxic bases include salts of primary,
- 20 secondary, and tertiary amines, substituted amines including naturally occurring substituted amines, cyclic amines and basic ion exchange resins, such as isopropylamine, trimethylamine, diethylamine, triethylamine, tripropylamine, and ethanolamine.

The pharmaceutical compositions can be prepared in various forms, such as granules,

25 tablets, pills, suppositories, capsules, suspensions, salves, lotions and the like. Pharmaceutical grade organic or inorganic carriers and/or diluents suitable for oral and topical use can be used to make up compositions containing the therapeutically-active compounds. Diluents known to the art include aqueous media, vegetable and animal oils and fats. Stabilizing agents, wetting and emulsifying agents, salts for

30 varying the osmotic pressure or buffers for securing an adequate pH value, and skin penetration enhancers can be used as auxiliary agents. The pharmaceutical

compositions may also include one or more of the following: carrier proteins such as serum albumin; buffers; fillers such as microcrystalline cellulose, lactose, corn and other starches; binding agents; sweeteners and other flavoring agents; coloring agents; and polyethylene glycol. Additives are well known in the art, and are used in a variety of formulations.

The administration of the mitotic agents of the present invention can be done in a variety of ways as discussed above, including, but not limited to, orally, subcutaneously, intravenously, intranasally, transdermally, intraperitoneally, intramuscularly, intrapulmonary, vaginally, rectally, or intraocularly. In some instances, for example, in the treatment of wounds and inflammation, the anti-mitotic agents may be directly applied as a solution or spray.

To employ the compounds of the invention in a method of screening for compounds that bind to KSP kinesin, the KSP is bound to a support, and a compound of the invention (which is a mitotic agent) is added to the assay. Alternatively, the compound of the invention is bound to the support and KSP is added. Classes of compounds among which novel binding agents may be sought include specific antibodies, non-natural binding agents identified in screens of chemical libraries, peptide analogs, etc. Of particular interest are screening assays for candidate agents that have a low toxicity for human cells. A wide variety of assays may be used for this purpose, including labeled in vitro protein-protein binding assays, electrophoretic mobility shift assays, immunoassays for protein binding, functional assays (phosphorylation assays, etc.) and the like.

The determination of the binding of the mitotic agent to KSP may be done in a number of ways. In a preferred embodiment, the mitotic agent (the compound of the invention) is labeled, for example, with a fluorescent or radioactive moiety and binding determined directly. For example, this may be done by attaching all or a portion of KSP to a solid support, adding a labeled mitotic agent (for example a compound of the invention in which at least one atom has been replaced by a detectable isotope), washing off excess reagent, and determining whether the amount of the label is that present on the solid support. Various blocking and washing steps may be utilized as is known in the art.

By "labeled" herein is meant that the compound is either directly or indirectly labeled with a label which provides a detectable signal, e.g., radioisotope, fluorescent tag, enzyme, antibodies, particles such as magnetic particles, chemiluminescent tag, or specific binding molecules, etc. Specific binding molecules include pairs, such as
5 biotin and streptavidin, digoxin and antidigoxin etc. For the specific binding members, the complementary member would normally be labeled with a molecule which provides for detection, in accordance with known procedures, as outlined above. The label can directly or indirectly provide a detectable signal.

In some embodiments, only one of the components is labeled. For example, the
10 kinesin proteins may be labeled at tyrosine positions using ^{125}I , or with fluorophores. Alternatively, more than one component may be labeled with different labels; using ^{125}I for the proteins, for example, and a fluorophor for the mitotic agents.

The compounds of the invention may also be used as competitors to screen for additional drug candidates. "Candidate bioactive agent" or "drug candidate" or
15 grammatical equivalents as used herein describe any molecule, e.g., protein, oligopeptide, small organic molecule, polysaccharide, polynucleotide, etc., to be tested for bioactivity. They may be capable of directly or indirectly altering the cellular proliferation phenotype or the expression of a cellular proliferation sequence, including both nucleic acid sequences and protein sequences. In other cases,
20 alteration of cellular proliferation protein binding and/or activity is screened. Screens of this sort may be performed either in the presence or absence of microtubules. In the case where protein binding or activity is screened, preferred embodiments exclude molecules already known to bind to that particular protein, for example, polymer structures such as microtubules, and energy sources such as ATP. Preferred
25 embodiments of assays herein include candidate agents which do not bind the cellular proliferation protein in its endogenous native state termed herein as "exogenous" agents. In another preferred embodiment, exogenous agents further exclude antibodies to KSP.

Candidate agents can encompass numerous chemical classes, though typically they are
30 organic molecules, preferably small organic compounds having a molecular weight of more than 100 and less than about 2,500 daltons. Candidate agents comprise

functional groups necessary for structural interaction with proteins, particularly hydrogen bonding and lipophilic binding, and typically include at least an amine, carbonyl, hydroxyl, ether, or carboxyl group, preferably at least two of the functional chemical groups. The candidate agents often comprise cyclical carbon or heterocyclic structures and/or aromatic or polyaromatic structures substituted with one or more of the above functional groups. Candidate agents are also found among biomolecules including peptides, saccharides, fatty acids, steroids, purines, pyrimidines, derivatives, structural analogs or combinations thereof. Particularly preferred are peptides.

Candidate agents are obtained from a wide variety of sources including libraries of synthetic or natural compounds. For example, numerous means are available for random and directed synthesis of a wide variety of organic compounds and biomolecules, including expression of randomized oligonucleotides. Alternatively, libraries of natural compounds in the form of bacterial, fungal, plant and animal extracts are available or readily produced. Additionally, natural or synthetically produced libraries and compounds are readily modified through conventional chemical, physical and biochemical means. Known pharmacological agents may be subjected to directed or random chemical modifications, such as acylation, alkylation, esterification, amidification to produce structural analogs.

Competitive screening assays may be done by combining KSP and a drug candidate in a first sample. A second sample comprises a mitotic agent, KSP and a drug candidate. This may be performed in either the presence or absence of microtubules. The binding of the drug candidate is determined for both samples, and a change, or difference in binding between the two samples indicates the presence of an agent capable of binding to KSP and potentially modulating its activity. That is, if the binding of the drug candidate is different in the second sample relative to the first sample, the drug candidate is capable of binding to KSP.

In a preferred embodiment, the binding of the candidate agent is determined through the use of competitive binding assays. In this embodiment, the competitor is a binding moiety known to bind to KSP, such as an antibody, peptide, binding partner, ligand, etc. Under certain circumstances, there may be competitive binding as

between the candidate agent and the binding moiety, with the binding moiety displacing the candidate agent.

In one embodiment, the candidate agent is labeled. Either the candidate agent, or the competitor, or both, is added first to KSP for a time sufficient to allow binding, if
5 present. Incubations may be performed at any temperature which facilitates optimal activity, typically between 4 and 40°C.

Incubation periods are selected for optimum activity, but may also be optimized to facilitate rapid high throughput screening. Typically between 0.1 and 1 hour will be sufficient. Excess reagent is generally removed or washed away. The second
10 component is then added, and the presence or absence of the labeled component is followed, to indicate binding.

In a preferred embodiment, the competitor is added first, followed by the candidate agent. Displacement of the competitor is an indication the candidate agent is binding to KSP and thus is capable of binding to, and potentially modulating, the activity of
15 KSP. In this embodiment, either component can be labeled. Thus, for example, if the competitor is labeled, the presence of label in the wash solution indicates displacement by the agent. Alternatively, if the candidate agent is labeled, the presence of the label on the support indicates displacement.

In an alternative embodiment, the candidate agent is added first, with incubation and
20 washing, followed by the competitor. The absence of binding by the competitor may indicate the candidate agent is bound to KSP with a higher affinity. Thus, if the candidate agent is labeled, the presence of the label on the support, coupled with a lack of competitor binding, may indicate the candidate agent is capable of binding to KSP.

25 It may be of value to identify the binding site of KSP. This can be done in a variety of ways. In one embodiment, once KSP has been identified as binding to the mitotic agent, KSP is fragmented or modified and the assays repeated to identify the necessary components for binding.

Modulation is tested by screening for candidate agents capable of modulating the activity of KSP comprising the steps of combining a candidate agent with KSP, as above, and determining an alteration in the biological activity of KSP. Thus, in this embodiment, the candidate agent should both bind to KSP (although this may not be
5 necessary), and alter its biological or biochemical activity as defined herein. The methods include both in vitro screening methods and in vivo screening of cells for alterations in cell cycle distribution, cell viability, or for the presence, morphology, activity, distribution, or amount of mitotic spindles, as are generally outlined above.

Alternatively, differential screening may be used to identify drug candidates that bind
10 to the native KSP, but cannot bind to modified KSP.

Positive controls and negative controls may be used in the assays. Preferably all control and test samples are performed in at least triplicate to obtain statistically significant results. Incubation of all samples is for a time sufficient for the binding of the agent to the protein. Following incubation, all samples are washed free of non-
15 specifically bound material and the amount of bound, generally labeled agent determined. For example, where a radiolabel is employed, the samples may be counted in a scintillation counter to determine the amount of bound compound.

A variety of other reagents may be included in the screening assays. These include reagents like salts, neutral proteins, e.g., albumin, detergents, etc which may be used
20 to facilitate optimal protein-protein binding and/or reduce non-specific or background interactions. Also reagents that otherwise improve the efficiency of the assay, such as protease inhibitors, nuclease inhibitors, anti-microbial agents, etc., may be used. The mixture of components may be added in any order that provides for the requisite binding.

25 The following examples serve to more fully describe the manner of using the above-described invention, as well as to set forth the best modes contemplated for carrying out various aspects of the invention. It is understood that these examples in no way serve to limit the true scope of this invention, but rather are presented for illustrative purposes. All references cited herein are incorporated by reference in their entirety.

EXAMPLES

Abbreviations and Definitions

The following abbreviations and terms have the indicated meanings throughout:

5	Ac	=	acetyl
	BNB	=	4-bromomethyl-3-nitrobenzoic acid
	Boc	=	t-butyloxy carbonyl
	Bu	=	butyl
	c-	=	cyclo
10	CBZ	=	carbobenzoxo = benzyloxycarbonyl
	DBU	=	diazabicyclo[5.4.0]undec-7-ene
	DCM	=	dichloromethane = methylene chloride = CH ₂ Cl ₂
	DCE	=	dichloroethylene
	DEAD	=	diethyl azodicarboxylate
15	DIC	=	diisopropylcarbodiimide
	DIEA	=	N,N-diisopropylethyl amine
	DMAP	=	4-N,N-dimethylaminopyridine
	DMF	=	N,N-dimethylformamide
	DMSO	=	dimethyl sulfoxide
20	DVB	=	1,4-divinylbenzene
	EEDQ	=	2-ethoxy-1-ethoxycarbonyl-1,2-dihydroquinoline
	Et	=	ethyl
	Fmoc	=	9-fluorenylmethoxycarbonyl
	GC	=	gas chromatography
25	HATU	=	O-(7-Azabenzotriazol-1-yl)-1,1,3,3-tetramethyluronium hexafluorophosphate
	HMDS	=	hexamethyldisilazane
	HOAc	=	acetic acid
	HOBt	=	hydroxybenzotriazole
30	Me	=	methyl
	mesyl	=	methanesulfonyl
	MTBE	=	methyl t-butyl ether
	NMO	=	N-methylmorpholine oxide
	PEG	=	polyethylene glycol
35	Ph	=	phenyl
	PhOH	=	phenol
	PfP	=	pentafluorophenol
	PPTS	=	pyridinium p-toluenesulfonate
	Py	=	pyridine
40	PyBroP	=	bromo-tris-pyrrolidino-phosphonium hexafluorophosphate
	rt	=	room temperature
	sat=d	=	saturated
	s-	=	secondary
	t-	=	tertiary
45	TBDMS	=	t-butyldimethylsilyl
	TES	=	triethylsilane

	TFA	=	trifluoroacetic acid
	THF	=	tetrahydrofuran
	TMOF	=	trimethyl orthoformate
	TMS	=	trimethylsilyl
5	tosyl	=	p-toluenesulfonyl
	Trt	=	triphenylmethyl

Example 1

Synthesis of Compounds

10 The general synthesis is shown in Figures 1 and 2.

Step 1: N-butyryl anthranilic acid.

To a three-necked, 500 mL round-bottom flask equipped with a thermometer, dropping funnel, and an efficient magnetic stir bar, was added anthranilic acid (1) (0.5 mole, 68.5 g) and dimethyl formamide (250 mL). To this solution was added butyryl
15 chloride (0.55 mole, 57.1 mL) dropwise at such a rate that the temperature of the mixture did not rise above 40°C. The suspension was stirred vigorously at room temperature for at least an additional 3 h. The mixture was poured into water (2000 mL) and stirred for another 1 h. The precipitated product was collected by filtration, washed with cold water, and dried under reduced pressure over P₂O₅, yielding
20 compound 2 (67.3 g, 65%).

Step 2: 2-Propyl-3,1-[4H]benzoxazin-4-one.

Compound 2 (51.8 g, 0.25 mole) was dissolved in acetic anhydride (180 mL) in a 500 mL round-bottom flask equipped with a magnetic stir bar, a Claisen-distillation head (with vacuum inlet) and a thermometer. The flask was placed in an oil bath and
25 slowly heated to 170-180°C with vigorous stirring. The acetic acid produced was slowly distilled off under atmospheric pressure. Monitoring the head temperature of the distillation unit was used to follow the progress of the transformation. The reaction mixture was then cooled to 60 °C and the excess of acetic anhydride removed by distillation under reduced pressure (ca. 20 mm Hg). The residue was afterward
30 cooled and the product crystallized. The product was triturated with n-hexane (75 mL) and isolated by filtration to yield 2-propyl-3,1-[4H]benzoxazin-4-one (3) (29.3 g, 62%). The above procedure gave compound 3 sufficiently pure to use directly in the

next step.

Step 3: 2-Propyl-3-benzylquinazolin-4-one.

Compound 3 (28.4 g, 0.15 mole) and benzylamine (17.5 mL, 0.16 mole) were refluxed in chloroform (50 ml) in a one-neck 250 mL round-bottom flask for 6 h.

- 5 After complete consumption of compound 3, the chloroform was evaporated under reduced pressure. Ethylene glycol (100 mL) and NaOH pellets (0.60 g) were added to the residue and the flask equipped with a Claisen-distillation head and a magnetic stir bar. The flask was immersed in an oil bath and reheated to 130-140 °C bath temperature with vigorous stirring and maintained there for 5 h while the water
10 produced was removed by distillation. After completion of the reaction, the clear solution was allowed to cool to room temperature and kept overnight to precipitate the product. The pH of the suspension was adjusted to 7-8 by adding 3% aq. HCl, the crystals were filtered off and washed with cold water, and then recrystallized from isopropanol (or alternatively from acetone) to provide the compound, 2-propyl-3-
15 benzylquinazolin-4-one (compound 4) (28.0 g, 67%).

Step 4: 2-(1'-bromopropyl)-3-benzylquinazolin-4-one.

- To a three-neck 250 mL round-bottom flask equipped with a thermometer, dropping funnel, and efficient magnetic stir bar was added compound 4 (27.8 g, 0.10 mole), anhydrous sodium acetate (10.0 g) and glacial acetic acid (130 mL). Bromine (16.0 g,
20 0.10 mole) dissolved in acetic acid (10 mL) was added dropwise to the above solution at 40 °C for 1-2 h. After addition was complete, the mixture was poured into water (1500 mL) and stirred for 1-2 h at room temperature. The precipitated product, 2-(1'-bromopropyl)-3-benzylquinazolin-4-one (5) was isolated by filtration, washed with warm water to remove traces of acetic acid, and rinsed with a small amount of
25 isopropanol. Drying yielded compound 5 (33.0 g, 92%).

Step 5: 2-[1'-(N,N-dimethylethylenediamino)propyl]-3-benzylquinazolin-4-one.

- Compound 5 (10.7 g, 0.03 mole) and N,N-dimethylethylenediamine (6.6 mL, 0.06 mole) were dissolved in abs. ethanol (60 mL) and heated at reflux for 6 h. After
30 completion of the reaction, the solvent was evaporated under reduced pressure. The


residue was dissolved in dichloromethane (150 mL) and washed with 3% aq. NaOH solution (ca. 10-20 mL). The organic layer was dried over MgSO_4 and evaporated to dryness under reduced pressure. The remaining oily product was purified by flash chromatography on a short silica gel pad using an eluent of CHCl_3 -MeOH-aq. NH_3 , 90:10:0.1, to give the desired compound (5), 2-[1'-(N,N-dimethylethylenediamino)propyl]-3-benzylquinazolin-4-one (6) (6.0 g, 55%).

Step 6: 2-[1'-(N-4-fluorobenzoyl)-(N,N-dimethylethylenediamino)propyl]-3-benzylquinazolin-4-one.

A stock solution of compound 5 (1.822 g, 5.0 mmol) was prepared in HPLC grade CHCl_3 (0.5 mL). A stock solution of p-fluorobenzoyl chloride (160.2 mg, 1 mmol) in HPLC grade 1,2-dichloroethane (2.0 mL) was prepared in a 2.0 mL volumetric flask. A third solution of triethylamine (2.0 mL of 0.5 M) was prepared in HPLC grade 1,2-dichloroethane. A 100 μL aliquot of each solution was pipetted into a glass reaction vessel using a Beckman Biomet 2000 automated liquid dispenser. The reaction mixture was shaken using a mechanical shaker, sonicated in an ultrasonic water bath, and then incubated overnight at room temperature. The mixture was diluted in CHCl_3 (300 μL) and washed with 5% aqueous NaHCO_3 and water. The solvent was removed in vacuo to provide compound 6 (65%). The purity of the compound was analyzed by TLC eluted with CH_2Cl_2 -ethanol-concentrated aqueous NH_3 , 100:10:1.

Examples 2 and 3

Synthesis of compounds of General Structure 1d



All anhydrous solvents were purchased from Aldrich chemical company in SureSeal®
 5 containers. Most reagents were purchase from Aldrich Chemical Company.
 Abbreviations: DCM, dichloromethane; DIEA, N,N-diisopropylethylamine; DMF,
 N,N-dimethylformamide; TES, triethylsilane; TFA, trifluoroacetic acid. Array
 synthesis was conducted in 15 x 75 mm glass round bottom screw-cap vials contained
 in a 4 x 6 array aluminum synthesis block, sealed with a Teflon-lined rubber
 10 membrane. Reagents were added and aqueous extractions performed with single or
 multichannel pipettors. Filtrations were performed using Whatman/Polyfiltronics 24
 well, 10 mL filtration blocks. Evaporation of volatile materials from the array was
 performed with a Labconco Vortex-Evaporator or by sweeping with a 4 x 6 nitrogen
 manifold.

15

Example 2 (solid phase synthesis of a single compound)

STEP 1) 1,3-Diaminopropane trityl resin (Novabiochem, 1.2 mmol/g) (0.20 g,
 0.24 mmol) was weighed into a screw-cap vial and 3 mL of a 1:1 mixture of DMF and
 chloroform was added. DIEA (0.130 mL, 0.72 mmol) and 2-(1'-bromopropyl)-3-
 20 benzylquinazolin-4-one (from Example 1) (0.188 g, 0.48 mmol) were added. The vial

was sealed, heated to 70 °C and shaken overnight. The resin was filtered and washed (3 x DCM, 2 x MeOH, 1 x DCM, 2 x ether) and dried under vacuum. A 27 mg aliquot of resin was treated with 5:5:90 TFA:TES:DCM for 15 min and the mixture was filtered and evaporated, resulting in 8 mg (64% yield) of the quinazolinone-diamine intermediate. LCMS analysis showed >80 % purity.

STEP 2) The resin from Step 1 was swelled in 3 mL of DCM. DIEA (0.130 mL, 0.72 mmol) and 4-bromobenzyl bromide (0.12 g, 0.48 mmol) were added. The vial was sealed and shaken overnight. LCMS analysis of a cleaved aliquot revealed an approximate 1:1 mixture of starting material and product. Another 0.130 mL of DIEA and 0.12 g of 4-bromobenzyl bromide were added and the mixture was shaken at 70 °C for 8 h. The resin was filtered, washed (as above), and dried under vacuum.

STEP 3) The resin from Step 2 was twice shaken for 30 min with 5:5:90 TFA:TES:DCM and filtered. The filtrates were combined and evaporated, yielding 140 mg of an orange oil. This material was purified by reverse phase preparative HPLC (acetonitrile-water gradient) to provide 27 mg (17% for 3 steps) of the mono-TFA salt.

Example 3 (combinatorial synthesis of multiple compounds)

STEP 1) 1,2-Diaminoethane trityl resin (Novabiochem, 0.95 mmol/g) (200 g, 1.9 mmol) and 1,3-Diaminopropane trityl resin (Novabiochem, 1.14 mmol/g) (2.0 g, 2.28 mmol) were each placed in different 10 mL polypropylene fritted tubes (Bio-Rad). To each were added 4 mL of DMF, 4 mL of chloroform, 3 eq. of DIEA (1.0 mL and 1.2 mL, respectively) and 2 eq. of 2-(1'-bromopropyl)-3-benzylquinazolin-4-one (from Example 1) (1.5 g and 1.8 g, respectively). The mixtures were shaken at 70 °C overnight. Each mixture was washed (3 x DCM, 2 x MeOH, 1 x DCM, 2 x ether) and dried under vacuum. Analysis of a cleaved aliquot revealed the presence of the appropriate quinazolinone-diamine for each in >90 % purity.

STEP 2) The quinazolinone ethyl-diamine resin (105 mg, 0.10 mmol) was placed into each of the vials in the first 2 rows of the array, and the quinazolinone propyl-diamine resin (88 mg, 0.10 mmol) was placed into each vial of the last 2 rows of the array. To each vial was added DIEA (0.131 mL, 0.75 mmol). Into each vial of the first 2 rows of the array was added a different amine, and the additions were repeated for the last two rows of the array. The reaction block was shaken at 70 °C


overnight. Liquid was removed from each vial by multichannel pipette using fine-pointed gel-well tips, and the resins were washed (2 x DCM, 1 x MeOH, 1 x DCM) and dried under vacuum.

- STEP 3) To each vial of the array was added 2 mL of a 10:5:85
5 TFA:TES:DCM solution. The reaction block was shaken for 45 min and the mixtures were transferred to a filter block, filtered, and washed twice with 0.75 mL DCM. The solutions were evaporated to yield yellow-to-red oils. These thick oils were triturated twice with ether, dissolved in DCM and treated with 4 M HCl in dioxane to provide
10 the HCl salts (unknown number of salts per compound) as tan-to-white powdery or amorphous solids. Analysis by LCMS showed all to be >75 % pure.

Examples 4-6


- Six racemic quinazolinones were separated into their enantiomers by chiral chromatography. The chiral chromatography of three of these compounds is
15 described below:

Example 4:



- Column - Chiralpak AD, 250 x 4.6 mm (Diacel Inc.). Sample - 0.5 mg/mL in EtOH.
Conditions - 15 min at 60% EtOH in Hexane, enantiomer 1 elutes at 4.5 min,
20 enantiomer 2 elutes at 4.9 min.

Example 5:




Column - Chiralcel OJ, 250 x 4.6 mm (Diacel Inc.). Sample – 0.5 mg/mL in EtOH.

Conditions - 15 min at 10% EtOH in Hexane, (R)-enantiomer elutes at 8.4 min, (S)-enantiomer elutes at 9.6 min.

5



Example 6:




Column - Chiralpak AD, 250 x 4.6 mm (Diacel Inc.). Sample – 0.5 mg/mL in EtOH.

10 Conditions - 15 min at 70% EtOH in Hexane, enantiomer 1 elutes at 6.5 min,
enantiomer 2 elutes at 8.8 min.



The table below depicts the IC₅₀ activity of the racemate and the enantiomers of three other compounds separated as above. In all three cases, one enantiomer was
15 significantly more potent than the other. By independent chiral synthesis, it appears
that the more active enantiomer is the R enantiomer.

	IC ₅₀ (μM) Racemate	IC ₅₀ (μM) Enantiomer 1	IC ₅₀ (μM) Enantiomer 2
 <chem>CC(C)N(CCCN(C)C)C(=O)c1ccc(Br)cc1C2=CN(Cc3ccccc3)C(=O)c4ccccc24</chem>	0.06	0.28	0.03
 <chem>CC(C)N(CCCN(C)C)C(=O)c1cccc2ccccc12C3=CN(Cc4ccccc4)C(=O)c5ccccc35</chem>	12.7	>>40	6.6

 <chem>CN(C)CCNC(=O)c1ccc(Br)cc1C2=C(N(C)C2c3ccccc3)c4ccccc4</chem>	2.6	>>40	1.3
---	-----	------	-----

Examples 7 and 8

The following two compounds were synthesized as single enantiomers by the route shown in Figure 4. The data indicate that the more active enantiomer is the R enantiomer.


	K_i (μ M)	K_i (μ M)
	S enantiomer	R enantiomer
	2	<0.1
	>0.5	<0.05

5

Example 9

Chiral Resolution by Recrystallization with Tartaric Acid

- 10 Intermediate A, prepared in Example 1, can be converted to an intermediate B, which, upon resolution, provides an alternative to the first five steps shown in Figure 4. The process is shown in the scheme below:




The *R* enantiomer of B can be crystallized selectively by heating a mixture of B with 1.1 equivalents of *D*-tartaric acid in a mixture of isopropanol and methanol and then
5 letting the mixture return to room temperature.

Example 9: X = Cl, R = H

Racemic intermediate B (1.5 g), dissolved in 100 mL of boiling isopropanol, was mixed with 0.8 g of *D*-tartaric acid in 100 mL of boiling methanol. The mixture was
10 allowed to slowly reach room temperature. After standing overnight, the solid was removed by filtration and rinsed with ethyl acetate and hexanes, and allowed to air dry. The dried solid (0.8 g) was then dissolved in a boiling mixture of 50 mL of isopropanol and 50 mL of methanol and allowed to slowly cool to room temperature. After standing overnight, the resulting solid was removed by filtration and rinsed with
15 ethyl acetate and hexanes, and allowed to air dry. The dried solid was then stirred with saturated sodium bicarbonate for 30 min and extracted with ethyl acetate. The organics were dried (MgSO₄), filtered and evaporated to dryness. The resulting clear oil weighed 345 mg. Chiral purity of >95% was determined by conversion of a portion to the S-Mosher amide and examination of the product by ¹HNMR.

20 The enantiomerically pure compounds below were prepared, according to the remaining steps in Figure 4, from material resulting from the procedure described above using both D- and L-tartaric acid.

	Racemic I C ₅₀ (uM)	R Isomer IC ₅₀ (uM)	S Isomer IC ₅₀ (uM)
	<0.05	<0.05	>0.5

Induction of Mitotic Arrest in Cell Populations Treated with a Quinazolinone KSP
Inhibitor

- 5 FACS analysis to determine cell cycle stage by measuring DNA content was performed as follows. Skov-3 cells (human ovarian cancer) were split 1:10 for plating in 10cm dishes and grown to subconfluence with RPMI 1640 medium containing 5% fetal bovine serum (FBS). The cells were then treated with either 10nM paclitaxel, 400nM quinazolinone 1, 200nM quinazolinone2, or 0.25% DMSO (vehicle for
- 10 compounds) for 24 hours. Cells were then rinsed off the plates with PBS containing 5mM EDTA, pelleted, washed once in PBS containing 1% FCS, and then fixed overnight in 85% ethanol at 4°C. Before analysis, the cells were pelleted, washed once, and stained in a solution of 10µg propidium iodide and 250µg of ribonuclease (RNase) A per milliliter at 37°C for half an hour. Flow cytometry analysis was
- 15 performed on a Becton-Dickinson FACScan, and data from 10,000 cells per sample was analyzed with Modfit software.

The quinazolinone compounds, as well as the known anti-mitotic agent paclitaxel, caused a shift in the population of cells from a G0/G1 cell cycle stage (2n DNA content) to a G2/M cell cycle stage (4n DNA content). Other compounds of this class were found to have similar effects.

5 Monopolar Spindle Formation following Application of a Quinazolinone KSP Inhibitor

To determine the nature of the G2/M accumulation, human tumor cell lines Skov-3 (ovarian), HeLa (cervical), and A549 (lung) were plated in 96-well plates at densities of 4,000 cells per well (SKOV-3 & HeLa) or 8,000 cells per well (A549), allowed to
10 adhere for 24 hours, and treated with various concentrations of the quinazolinone compounds for 24 hours. Cells were fixed in 4% formaldehyde and stained with anti-tubulin antibodies (subsequently recognized using fluorescently-labeled secondary antibody) and Hoechst dye (which stains DNA).

Visual inspection revealed that the quinazolinone compounds caused cell cycle arrest in the prometaphase stage of mitosis. DNA was condensed and spindle formation had initiated, but arrested cells uniformly displayed monopolar spindles, indicating that there was an inhibition of spindle pole body separation. Microinjection of anti-KSP antibodies also causes mitotic arrest with arrested cells displaying monopolar spindles.

Inhibition of Cellular Proliferation in Tumor Cell Lines Treated with Quinazolinone
KSP Inhibitors.

Cells were plated in 96-well plates at densities from 1000-2500 cells/well of a 96-well plate (depending on the cell line) and allowed to adhere/grow for 24 hours. They were then treated with various concentrations of drug for 48 hours. The time at which compounds are added is considered T_0 . A tetrazolium-based assay using the reagent 3-(4,5-dimethylthiazol-2-yl)-5-(3-carboxymethoxyphenyl)-2-(4-sulfophenyl)-2H-tetrazolium (MTS) (I.S> Patent No. 5,185,450) (see Promega product catalog #G3580, CellTiter 96® AQueous One Solution Cell Proliferation Assay) was used to determine the number of viable cells at T_0 and the number of cells remaining after 48 hours compound exposure. The number of cells remaining after 48 hours was compared to

the number of viable cells at the time of drug addition, allowing for calculation of growth inhibition.

The growth over 48 hours of cells in control wells that had been treated with vehicle only (0.25% DMSO) is considered 100% growth and the growth of cells in wells with
5 compounds is compared to this. Quinazolinone KSP inhibitors inhibited cell proliferation in human tumor cell lines of the following tumor types: lung (NCI-H460, A549), breast (MDA-MB-231, MCF-7, MCF-7/ADR-RES), colon (HT29, HCT15), ovarian (SKOV-3, OVCAR-3), leukemia (HL-60(TB), K-562), central nervous system (SF-268), renal (A498), osteosarcoma (U2-OS), and cervical (HeLa).
10 In addition, a mouse tumor line (B16, melanoma) was also growth-inhibited in the presence of the quinazolinone compounds.

A Gi_{50} was calculated by plotting the concentration of compound in μM vs the percentage of cell growth of cell growth in treated wells. The Gi_{50} calculated for the compounds is the estimated concentration at which growth is inhibited by 50%
15 compared to control, i.e., the concentration at which:

$$100 \times [(Treated_{48} - T_0) / (Control_{48} - T_0)] = 50.$$

All concentrations of compounds are tested in duplicate and controls are averaged over 12 wells. A very similar 96-well plate layout and Gi_{50} calculation scheme is used by the National Cancer Institute (see Monks, et al., J. Natl. Cancer Inst. 83:757-766
20 (1991)). However, the method by which the National Cancer Institute quantitates cell number does not use MTS, but instead employs alternative methods.

Calculation Of IC_{50} :

Measurement of a composition's IC_{50} for KSP activity uses an ATPase assay. The following solutions are used: Solution 1 consists of 3 mM phosphoenolpyruvate
25 potassium salt (Sigma P-7127), 2 mM ATP (Sigma A-3377), 1 mM IDTT (Sigma D-9779), 5 μM paclitaxel (Sigma T-7402), 10 ppm antifoam 289 (Sigma A-8436), 25 mM Pipes/KOH pH 6.8 (Sigma P6757), 2 mM $MgCl_2$ (VWR JT400301), and 1 mM EGTA (Sigma E3889). Solution 2 consists of 1 mM NADH (Sigma N8129), 0.2 mg/ml BSA (Sigma A7906), pyruvate kinase 7U/ml, L-lactate dehydrogenase 10 U/ml

(Sigma P0294), 100 nM KSP motor domain, 50 µg/ml microtubules, 1 mM DTT (Sigma D9779), 5 µM paclitaxel (Sigma T-7402), 10 ppm antifoam 289 (Sigma A-8436), 25 mM Pipes/KOH pH 6.8 (Sigma P6757), 2 mM MgCl₂ (VWR JT4003-01), and 1 mM EGTA (Sigma E3889). Serial dilutions (8-12 two-fold dilutions) of the composition are made in a 96-well microtiter plate (Corning Costar 3695) using Solution 1. Following serial dilution each well has 50 µl of Solution 1. The reaction is started by adding 50 µl of solution 2 to each well. This may be done with a multichannel pipettor either manually or with automated liquid handling devices. The microtiter plate is then transferred to a microplate absorbance reader and multiple absorbance readings at 340 nm are taken for each well in a kinetic mode. The observed rate of change, which is proportional to the ATPase rate, is then plotted as a function of the compound concentration. For a standard IC₅₀ determination the data acquired is fit by the following four parameter equation using a nonlinear fitting program (e.g., Grafit 4):

$$y = \frac{\text{Range}}{1 + \left(\frac{x}{\text{IC}_{50}} \right)^s} + \text{Background}$$

Where y is the observed rate and x the compound concentration.



The quinazolinone compounds inhibit growth in a variety of cell lines, including cell lines (MCF-7/ADR-RES, HCT1 5) that express P-glycoprotein (also known as Multi-drug Resistance, or MDR⁺), which conveys resistance to other chemotherapeutic drugs, such as paclitaxel. Therefore, the quinazolinones are anti-mitotics that inhibit cell proliferation, and are not subject to resistance by overexpression of MDR⁺ by drug-resistant tumor lines.

Other compounds of this class were found to inhibit cell proliferation, although GI₅₀ values varied. GI₅₀ values for the quinazolinone compounds tested ranged from 200 nM to greater than the highest concentration tested. By this we mean that although most of the compounds that inhibited KSP activity biochemically did inhibit cell proliferation, for some, at the highest concentration tested (generally about 20 µM), cell growth was inhibited less than 50%. Many of the compounds have GI₅₀



values less than 10 μM , and several have GI_{50} values less than 1 μM . Anti-proliferative compounds that have been successfully applied in the clinic to treatment of cancer (cancer chemotherapeutics) have GI_{50} 's that vary greatly. For example, in A549 cells, paclitaxel GI_{50} is 4 nM, doxorubicin is 63 nM, 5-fluorouracil is 1 μM , and
5 hydroxyurea is 500 μM (data provided by National Cancer Institute, Developmental Therapeutic Program, <http://dtp.nci.nih.gov/>). Therefore, compounds that inhibit cellular proliferation at virtually any concentration may be useful. However, preferably, compounds will have GI_{50} values of less than 1 mM. More preferably, compounds will have GI_{50} values of less than 20 μM . Even more preferably,
10 compounds will have GI_{50} values of less than 10 μM . Further reduction in GI_{50} values may also be desirable, including compounds with GI_{50} values of less than 1 μM . Some of the quinazolinone compounds of the invention inhibit cell proliferation with GI_{50} values from below 200 nM to below 10 nM.

We claim:

1. A method of treating cellular proliferative diseases comprising administering a compound chosen from the group consisting of:



5



and

wherein:

R_1 is chosen from hydrogen, alkyl, aryl, alkylaryl, heteroaryl, alkylheteroaryl, substituted alkyl, substituted aryl, substituted alkylaryl, substituted heteroaryl, and substituted alkylheteroaryl;

10

R_2 and R_2' are independently chosen from hydrogen, alkyl, oxaalkyl, aryl, alkylaryl, heteroaryl, alkylheteroaryl, substituted alkyl, substituted aryl, substituted alkylaryl, substituted heteroaryl, and substituted alkylheteroaryl; or R_2 and R_2' taken together form a 3- to 7-membered ring;

15

R_3 is chosen from hydrogen, alkyl, aryl, alkylaryl, heteroaryl, alkylheteroaryl, substituted alkyl, substituted aryl, substituted alkylaryl, substituted heteroaryl, substituted alkylheteroaryl, oxaalkyl, oxaalkylaryl, substituted oxaalkylaryl, $R_{15}O-$ and $R_{15}NH-$;

R_3 is chosen from hydrogen, alkyl, aryl, alkylaryl, heteroaryl, alkylheteroaryl, substituted alkyl, substituted aryl, substituted alkylaryl, substituted heteroaryl, substituted alkylheteroaryl, and R_{15} -NH-;

5 $R_{3'}$ is chosen from alkyl, aryl, alkylaryl, heteroaryl, alkylheteroaryl, substituted alkyl, substituted aryl, substituted alkylaryl, substituted heteroaryl, and substituted alkylheteroaryl;



R_4 is chosen from hydrogen, alkyl, aryl, alkylaryl, heteroaryl, alkylheteroaryl, substituted alkyl, substituted aryl, substituted alkylaryl, substituted heteroaryl, substituted alkylheteroaryl, and R_{16} -alkylene-;


10 R_5 , R_6 , R_7 and R_8 are independently chosen from hydrogen, alkyl, alkoxy, halogen, fluoroalkyl, nitro, dialkylamino, alkylsulfonyl, alkylsulfonamido, sulfonamidoalkyl, sulfonamidoaryl, alkylthio, carboxyalkyl, carboxamido, aminocarbonyl, aryl and heteroaryl;

15 R_{15} is chosen from alkyl, aryl, alkylaryl, heteroaryl, alkylheteroaryl, substituted alkyl, substituted aryl, substituted alkylaryl, substituted heteroaryl, and substituted alkylheteroaryl;

R_{16} is chosen from alkoxy, amino, alkylamino, dialkylamino, N-heterocyclyl and substituted N-heterocyclyl.

2. A method of treating a disorder associated with KSP kinesin activity
20 comprising administering a compound chosen from the group consisting of:





wherein:



- R_1 is chosen from hydrogen, alkyl, aryl, alkylaryl, heteroaryl, alkylheteroaryl, substituted alkyl, substituted aryl, substituted alkylaryl, substituted heteroaryl, and substituted alkylheteroaryl;
- R_2 and R_2' are independently chosen from hydrogen, alkyl, oxaalkyl, aryl, alkylaryl, heteroaryl, alkylheteroaryl, substituted alkyl, substituted aryl, substituted alkylaryl, substituted heteroaryl, and substituted alkylheteroaryl; or R_2 and R_2' taken together form a 3- to 7-membered ring;
- R_3 is chosen from hydrogen, alkyl, aryl, alkylaryl, heteroaryl, alkylheteroaryl, substituted alkyl, substituted aryl, substituted alkylaryl, substituted heteroaryl, substituted alkylheteroaryl, oxaalkyl, oxaalkylaryl, substituted oxaalkylaryl, $R_{15}O$ - and $R_{15}NH$ -;
- R_3' is chosen from hydrogen, alkyl, aryl, alkylaryl, heteroaryl, alkylheteroaryl, substituted alkyl, substituted aryl, substituted alkylaryl, substituted heteroaryl, substituted alkylheteroaryl and $R_{15}NH$ -;
- R_3'' is chosen from alkyl, aryl, alkylaryl, heteroaryl, alkylheteroaryl, substituted alkyl, substituted aryl, substituted alkylaryl, substituted heteroaryl, and substituted alkylheteroaryl;
- R_4 is chosen from hydrogen, alkyl, aryl, alkylaryl, heteroaryl, alkylheteroaryl, substituted alkyl, substituted aryl, substituted alkylaryl, substituted heteroaryl, substituted alkylheteroaryl, and R_{16} -alkylene-;
- R_5 , R_6 , R_7 and R_8 are independently chosen from hydrogen, alkyl, alkoxy, halogen, fluoroalkyl, nitro, dialkylamino, alkylsulfonyl, alkylsulfonamido, sulfonamidoalkyl, sulfonamidoaryl, alkylthio, carboxyalkyl, carboxamido,

aminocarbonyl, aryl and heteroaryl;



R_{15} is chosen from alkyl, aryl, alkylaryl, heteroaryl, alkylheteroaryl, substituted alkyl, substituted aryl, substituted alkylaryl, substituted heteroaryl, and substituted alkylheteroaryl;

- 5 R_{16} is chosen from alkoxy, amino, alkylamino, dialkylamino, N-heterocyclyl and substituted N-heterocyclyl.

3. A method of inhibiting KSP kinesin comprising contacting KSP kinesin with a compound chosen from the group consisting of:



10



and

wherein:

R_1 is chosen from hydrogen, alkyl, aryl, alkylaryl, heteroaryl, alkylheteroaryl, substituted alkyl, substituted aryl, substituted alkylaryl, substituted heteroaryl, and substituted alkylheteroaryl;

15

R_2 and $R_{2'}$ are independently chosen from hydrogen, alkyl, oxaalkyl, aryl, alkylaryl, heteroaryl, alkylheteroaryl, substituted alkyl, substituted aryl, substituted alkylaryl, substituted heteroaryl, and substituted alkylheteroaryl; or R_2 and $R_{2'}$ taken together form a 3- to 7-membered ring;

R₃ is chosen from hydrogen, alkyl, aryl, alkylaryl, heteroaryl, alkylheteroaryl, substituted alkyl, substituted aryl, substituted alkylaryl, substituted heteroaryl, substituted alkylheteroaryl, oxaalkyl, oxaalkylaryl, substituted oxaalkylaryl, R₁₅O- and R₁₅-NH-;

- 5 R_{3'} is chosen from hydrogen, alkyl, aryl, alkylaryl, heteroaryl, alkylheteroaryl, substituted alkyl, substituted aryl, substituted alkylaryl, substituted heteroaryl, substituted alkylheteroaryl and R₁₅-NH-;

- R_{3''} is chosen from alkyl, aryl, alkylaryl, heteroaryl, alkylheteroaryl, substituted alkyl, substituted aryl, substituted alkylaryl, substituted heteroaryl, and substituted
10 alkylheteroaryl;

R₄ is chosen from hydrogen, alkyl, aryl, alkylaryl, heteroaryl, alkylheteroaryl, substituted alkyl, substituted aryl, substituted alkylaryl, substituted heteroaryl, substituted alkylheteroaryl, and R₁₆-alkylene-;

- R₅, R₆, R₇ and R₈ are independently chosen from hydrogen, alkyl, alkoxy, halogen,
15 fluoroalkyl, nitro, dialkylamino, alkylsulfonyl, alkylsulfonamido, sulfonamidoalkyl, sulfonamidoaryl, alkylthio, carboxyalkyl, carboxamido, aminocarbonyl, aryl and heteroaryl;


- R₁₅ is chosen from alkyl, aryl, alkylaryl, heteroaryl, alkylheteroaryl, substituted alkyl, substituted aryl, substituted alkylaryl, substituted heteroaryl, and substituted
20 alkylheteroaryl;

R₁₆ is chosen from alkoxy, amino, alkylamino, dialkylamino, N-heterocyclyl and substituted N-heterocyclyl.

4. A method according to claim 1, 2 or 3 wherein
- R_1 is chosen from hydrogen, alkyl, aryl, substituted alkyl, substituted aryl, heteroaryl, substituted heteroaryl, alkylaryl, substituted alkylaryl and substituted alkylheteroaryl;
- 5 R_2 is chosen from hydrogen, alkyl and substituted alkyl;
 R_2' is hydrogen;
- R_3 is chosen from alkyl, substituted alkyl, alkylaryl, heteroaryl, aryl, substituted aryl, substituted heteroaryl, substituted oxaalkylaryl $R_{15}O-$ and $R_{15}NH-$;
- R_4 is chosen from alkyl, aryl, alkylaryl, alkylheteroaryl, substituted alkyl, substituted
- 10 aryl, and R_{16} -alkylene-;
- R_5 is hydrogen;
- R_6, R_7 and R_8 are independently chosen from hydrogen, halogen, methyl and trifluoromethyl;
- R_{15} is chosen from alkyl, aryl and substituted aryl;
- 15 R_{16} is chosen from alkoxy, amino, alkylamino, dialkylamino and N-heterocyclyl.

5. A method according to claim 4 wherein the stereogenic center to which R_2 and R_2' are attached is of the R configuration.

- 20 6. A method according to claim 1, 2 or 3 comprising administering a compound of formula:



7. A method according to claim 6 wherein R_1 is chosen from hydrogen, lower alkyl, substituted lower alkyl, benzyl, substituted benzyl, phenyl, naphthyl and substituted phenyl.
8. A method according to claim 7 wherein R_1 is chosen from hydrogen,
5 ethyl, propyl, methoxyethyl, naphthyl, phenyl, bromophenyl, chlorophenyl, methoxyphenyl, ethoxyphenyl, tolyl, dimethylphenyl, chlorofluorophenyl, methylchlorophenyl, ethylphenyl, phenethyl, benzyl, chlorobenzyl, methylbenzyl, methoxybenzyl, tetrahydrofuranylmethyl and (ethoxycarbonyl)ethyl.
9. A method according to claim 6 wherein R_2 is chosen from hydrogen,
10 lower alkyl and substituted lower alkyl, and R_2' is hydrogen.
10. A method according to claim 9 wherein R_2 is chosen from hydrogen, methyl, ethyl, propyl, methylthioethyl, aminobutyl, (CBZ)aminobutyl, cyclohexylmethyl, benzyloxymethyl, methylsulfinylethyl, methylsulfinylmethyl, hydroxymethyl, benzyl and indolylmethyl.
11. A method according to claim 6 wherein R_3 is chosen from C_1 - C_{13}
15 alkyl; substituted lower alkyl; phenyl; naphthyl; phenyl substituted with one or more halo, lower alkyl, loweralkoxy, nitro, carboxy, methylenedioxy or trifluoromethyl; biphenyl; benzyl; phenoxymethyl; halophenoxymethyl; phenylvinyl; heteroaryl; heteroaryl substituted with lower alkyl; and benzyloxymethyl.
12. A method according to claim 11 wherein R_3 is chosen from ethyl,
20 propyl, chloropropyl, butoxy, heptyl, butyl, octyl, tridecanyl, (ethoxycarbonyl)ethyl, dimethylaminoethyl, dimethylaminomethyl, phenyl, naphthyl, halophenyl, dihalophenyl, cyanophenyl, halo(trifluoromethyl)phenyl, chlorophenoxymethyl, methoxyphenyl, carboxyphenyl, ethylphenyl, tolyl, biphenyl, methylenedioxyphenyl,
25 methylsulfonylphenyl, methoxychlorophenyl, chloronaphthyl, methylhalophenyl, trifluoromethylphenyl, butylphenyl, pentylphenyl, methylnitrophenyl, phenoxymethyl, dimethoxyphenyl, phenylvinyl, nitrochlorophenyl, nitrophenyl, dinitrophenyl, bis(trifluoromethyl)phenyl, benzyloxymethyl, benzyl, furanyl, benzofuranyl, pyridinyl,

indolyl, methylpyridinyl, quinolinyl, picolinyl, pyrazolyl, and imidazolyl.

13. A method according to claim 6 wherein R_3 is R_{15} -NH- and R_{15} is chosen from lower alkyl; cyclohexyl; phenyl; and phenyl substituted with halo, lower alkyl, loweralkoxy, or lower alkylthio.

5 14. A method according to claim 13 wherein R_{15} is chosen from isopropyl, butyl, cyclohexyl, phenyl, bromophenyl, dichlorophenyl, methoxyphenyl, ethylphenyl, tolyl, trifluoromethylphenyl and methylthiophenyl.

15. A method according to claim 6 wherein R_4 is chosen from lower alkyl, substituted lower alkyl, cyclohexyl; phenyl substituted with hydroxy, lower alkoxy or
10 lower alkyl; benzyl; heteroarylmethyl; heteroarylethyl; heteroarylpropyl and R_{16} -alkylene-, wherein R_{16} is amino, lower alkylamino, di(lower alkyl)amino, lower alkoxy, or N-heterocyclyl.

16. A method according to claim 15 wherein R_4 is chosen from methyl, ethyl, propyl, butyl, cyclohexyl, carboxyethyl, carboxymethyl, methoxyethyl,
15 hydroxyethyl, hydroxypropyl, dimethylaminoethyl, dimethylaminopropyl, diethylaminoethyl, diethylaminopropyl, aminopropyl, methylaminopropyl, 2,2-dimethyl-3-(dimethylamino)propyl, 1-cyclohexyl-4-(diethylamino)butyl, aminoethyl, aminobutyl, aminopentyl, aminohexyl, aminoethoxyethyl, isopropylaminopropyl, diisopropylaminoethyl, 1-methyl-4-(diethylamino)butyl, (t-Boc)aminopropyl,
20 hydroxyphenyl, benzyl, methoxyphenyl, methylmethoxyphenyl, dimethylphenyl, tolyl, ethylphenyl, (oxopyrrolidinyl)propyl, (methoxycarbonyl)ethyl, benzylpiperidinyl, pyridinylethyl, pyridinylmethyl, morpholinylethyl, morpholinylpropyl, piperidinyl, azetidinylmethyl, azetidinypropyl, pyrrolidinylethyl, pyrrolidinylpropyl, piperidinylmethyl, piperidinylethyl, imidazolylpropyl, imidazolylethyl,
25 (ethylpyrrolidinyl)methyl, (methylpyrrolidinyl)ethyl, (methylpiperidinyl)propyl, (methylpiperazinyl)propyl, furanylmethyl and indolylethyl.

17. A method according to claim 6 wherein R_1 is chosen from lower alkyl, benzyl, substituted benzyl and substituted phenyl;

R₂ is chosen from hydrogen, alkyl, substituted lower alkyl and benzyl;

R₂' is hydrogen;

R₃ is chosen from substituted phenyl and naphthyl;

R₄ is chosen from substituted alkyl and R₁₆-alkylene-;

5 R₅ is hydrogen or halo

R₆ is hydrogen, methyl or halo;

R₇ is hydrogen, halo, methyl or trifluoromethyl;

R₈ is hydrogen or halo;

R₁₆ is chosen from di(lower alkylamino), (lower alkyl)amino, amino, N-heterocyclyl

10 and substituted N-heterocyclyl.

18. A method according to claim 1, 2 or 3 wherein

R₁ is benzyl or halobenzyl;

R₂ is chosen from ethyl and propyl;

15 R₂' is hydrogen;

R₃ is substituted phenyl;

R₃' is substituted phenyl;

R₃'' is substituted phenyl;

R₄ is (CH₂)_m OH or (CH₂)_p R₁₆ wherein m is 2 or 3 and p is 1-3;

20 R₅ is hydrogen;

R₆ is hydrogen;

R₇ is halo;


R₈ is hydrogen;

R₁₆ is chosen from amino, propylamino, and azetidiny.

25

19. A method according to claim 18 wherein the stereogenic center to which R₂ and R₂' are attached is of the R configuration.

20. A method according to claim 1, 2 or 3 comprising administering a
30 compound of formula:



21. A method according to claim 20 wherein:

R₁ is chosen from hydrogen, lower alkyl, substituted lower alkyl, benzyl, substituted benzyl, phenyl, naphthyl and substituted phenyl;

5 R₂ is chosen from hydrogen, lower alkyl and substituted lower alkyl and R₂' is hydrogen;

R₃' is chosen from C₁-C₁₃ alkyl; phenyl; naphthyl; phenyl substituted with halo, lower alkyl, lower alkoxy, nitro, methylenedioxy, or trifluoromethyl; biphenyl, benzyl and heteroaryl; and

10 R₄ is chosen from lower alkyl, substituted lower alkyl, cyclohexyl; phenyl substituted with hydroxy, lower alkoxy or lower alkyl; benzyl; heteroarylmethyl; heteroarylethyl; heteroarylpropyl and R₁₆-alkylene, wherein

R₁₆ is amino, (lower alkyl)amino, di(lower alkyl)amino, lower alkoxy, or N-heterocyclyl.

15 22. A method according to claim 20 wherein

R₁ is chosen from lower alkyl, benzyl, substituted benzyl and substituted phenyl;

R₂ is hydrogen or lower alkyl;

R₂' is hydrogen;

R₃ is chosen from substituted phenyl and naphthyl;


20 R₄ is R₁₆-alkylene- hydroxy lower alkyl or carboxy lower alkyl;

R₆ and R₇ are chosen from hydrogen and halo;

R₅ and R₈ are hydrogen;

R₁₆ is chosen from di(lower alkylamino), (lower alkyl)amino, amino, piperidinyl, azetidiny, pyrrolidinyl and morpholinyl.

23. A method according to claim 1, 2 or 3 comprising administering a compound of formula:



24. A method according to claim 23 wherein:

5 R₁ is chosen from hydrogen, lower alkyl, substituted lower alkyl, benzyl, substituted benzyl, phenyl, naphthyl and substituted phenyl;

R₂ is chosen from hydrogen, lower alkyl and substituted lower alkyl and R₂' is hydrogen; and

10 R₄ is chosen from lower alkyl, cyclohexyl; phenyl substituted with hydroxy, lower alkoxy or lower alkyl; benzyl; heteroarylmethyl; heteroarylethyl; heteroarylpropyl and R₁₆-alkylene, wherein R₁₆ is di(lower alkyl)amino, alkylamino, amino, lower alkoxy, or N-heterocyclyl.

25. A method according to claim 23 wherein

R₁ is chosen from lower alkyl, benzyl, substituted benzyl and substituted phenyl;

15 R₂ is hydrogen or lower alkyl;

R₂' is hydrogen;

R₄ is R₁₆-alkylene-;


R₆ and R₇ are chosen from hydrogen and halo;

R₅ and R₈ are hydrogen;

20 R₁₆ is chosen from di(lower alkylamino), (lower alkyl)amino, amino, pyrrolidinyl, piperidinyl, imidazolyl and morpholinyl.

26. A method according to claim 1, 2 or 3 comprising administering a

compound of formula:



27. A method according to claim 26 wherein:

- R₁ is chosen from hydrogen, lower alkyl, substituted lower alkyl, benzyl, substituted
 5 benzyl, phenyl, naphthyl and substituted phenyl;
 R₂ is chosen from hydrogen, lower alkyl and substituted lower alkyl and R₂' is
 hydrogen;
 R₃'' is chosen from C₁-C₁₃ alkyl; substituted lower alkyl; phenyl; naphthyl; phenyl
 substituted with halo, lower alkyl, lower alkoxy, nitro, methylenedioxy, or
 10 trifluoromethyl; biphenyl; benzyl and heterocyclyl; and
 R₄ is chosen from lower alkyl, substituted lower alkyl; cyclohexyl; phenyl substituted
 with hydroxy, lower alkoxy or lower alkyl; benzyl; substituted benzyl;
 heterocyclyl; heteroarylmethyl; heteroarylethyl; heteroarylpropyl and R₁₆-
 alkylene, wherein
 15 R₁₆ is di(lower alkyl)amino, (lower alkyl)amino, amino, lower alkoxy, or N-
 heterocyclyl.

28. A method according to claim 27 wherein


- R₁ is chosen from lower alkyl, benzyl, substituted benzyl and substituted phenyl;
R₂ is hydrogen or lower alkyl;
R₂' is hydrogen;
5 R₃'' is chosen from substituted phenyl, heterocyclyl and naphthyl;
R₄ is chosen from substituted benzyl, heterocyclyl substituted lower alkyl and R₁₆-alkylene-;
R₆ and R₇ are chosen from hydrogen and halo;
R₅ and R₈ are hydrogen;
10 R₁₆ is chosen from di(lower alkylamino), (lower alkyl)amino, amino, pyrrolidinyl, azetidiny, piperidinyl, imidazolyl and morpholinyl.

29. A method according to claim 28 wherein

- R₁ is benzyl;
R₂ is ethyl;
15 R₂' is hydrogen;
R₃'' is chosen from halophenyl, polyhalophenyl, tolyl, dimethylphenyl, methoxyphenyl, dimethoxyphenyl, cyanophenyl, trifluoromethylphenyl, trifluoromethoxyphenyl, bis(trifluoromethyl)phenyl, carboxyphenyl, t-butylphenyl, methoxycarbonylphenyl, piperidinyl and naphthyl;
20 R₄ is chosen from substituted benzyl, piperidinyl, hydroxy (lower alkyl) and R₁₆-alkylene-;
R₆ and R₇ are chosen from hydrogen and halo;
R₅ and R₈ are hydrogen;
R₁₆ is chosen from dimethylamino, amino, pyrrolidinyl and piperidinyl.

- 25 30. A method according to claim 1 or 2 wherein said disease or disorder is chosen from the group consisting of cancer, hyperplasia, restenosis, cardiac hypertrophy, immune disorders and inflammation.

31. A compound chosen from the group consisting of:



5

wherein:

R_1 is chosen from hydrogen, alkyl, aryl, alkylaryl, heteroaryl, alkylheteroaryl, substituted alkyl, substituted aryl, substituted alkylaryl, substituted heteroaryl, and substituted alkylheteroaryl;

10 R_2 and R_2' are independently chosen from hydrogen, alkyl, oxaalkyl, aryl, alkylaryl, heteroaryl, alkylheteroaryl, substituted alkyl, substituted aryl, substituted alkylaryl, substituted heteroaryl, and substituted alkylheteroaryl; or R_2 and R_2' taken together form a 3- to 7-membered ring;

R_3 is chosen from hydrogen, alkyl, aryl, alkylaryl, heteroaryl, alkylheteroaryl, substituted alkyl, substituted aryl, substituted alkylaryl, substituted heteroaryl, substituted alkylheteroaryl, oxaalkyl, oxaalkylaryl, substituted oxaalkylaryl, $R_{15}O-$ and $R_{15}NH-$;

15

R₃ is chosen from hydrogen, alkyl, aryl, alkylaryl, heteroaryl, alkylheteroaryl, substituted alkyl, substituted aryl, substituted alkylaryl, substituted heteroaryl, substituted alkylheteroaryl and R₁₅-NH-;

5 R₃' is chosen from alkyl, aryl, alkylaryl, heteroaryl, alkylheteroaryl, substituted alkyl, substituted aryl, substituted alkylaryl, substituted heteroaryl, and substituted alkylheteroaryl;

R₄ is chosen from hydrogen, alkyl, aryl, alkylaryl, heteroaryl, alkylheteroaryl, substituted alkyl, substituted aryl, substituted alkylaryl, substituted heteroaryl, substituted alkylheteroaryl, and R₁₆-alkylene-;

10 R₅, R₆, R₇ and R₈ are independently chosen from hydrogen, alkyl, alkoxy, halogen, fluoroalkyl, nitro, dialkylamino, alkylsulfonyl, alkylsulfonamido, sulfonamidoalkyl, sulfonamidoaryl, alkylthio, carboxyalkyl, carboxamido, aminocarbonyl, aryl and heteroaryl;

15 R₁₅ is chosen from hydrogen, alkyl, aryl, alkylaryl, heteroaryl, alkylheteroaryl, substituted alkyl, substituted aryl, substituted alkylaryl, substituted heteroaryl, and substituted alkylheteroaryl;

R₁₆ is chosen from alkoxy, amino, alkylamino, dialkylamino, N-heterocyclyl and substituted N-heterocyclyl;

20 with the proviso that when R₃ is R₁₅-NH- attached to carbonyl, both of R₂ and R₄ must be other than hydrogen.

32. A compound according to claim 31 wherein

R₁ is chosen from hydrogen, alkyl, aryl, substituted alkyl, substituted aryl, heteroaryl, substituted heteroaryl, alkylaryl, alkylheteroaryl and substituted alkylaryl;

R₂ is chosen from hydrogen, alkyl and substituted alkyl;

25 R₂' is hydrogen;

R₃ is chosen from alkyl, aryl, alkylaryl, heteroaryl, substituted aryl, substituted alkyl, substituted heteroaryl, oxaalkylaryl, substituted oxaalkylaryl, R₁₅O- and R₁₅-NH-;

30 R₄ is chosen from alkyl, aryl, alkylaryl, alkylheteroaryl, substituted alkyl, substituted aryl, and R₁₆-alkylene-;


R₅ is hydrogen;

R₆, R₇ and R₈ are independently chosen from hydrogen, halogen, methyl and trifluoromethyl;

R₁₅ is chosen from alkyl, aryl and substituted aryl;

5 R₁₆ is chosen from alkoxy, amino, alkylamino, dialkylamino and N-heterocyclyl.

33. A compound according to claim 31 of formula:



34. A compound according to claim 33 wherein R₁ is chosen from hydrogen, lower alkyl, substituted lower alkyl, benzyl, substituted benzyl, phenyl,
10 naphthyl and substituted phenyl.

35. A compound according to claim 34 wherein R₁ is chosen from hydrogen, ethyl, propyl, methoxyethyl, naphthyl, phenyl, bromophenyl, chlorophenyl, methoxyphenyl, ethoxyphenyl, tolyl, dimethylphenyl, chlorofluorophenyl, methylchlorophenyl, ethylphenyl, phenethyl, benzyl, chlorobenzyl, methylbenzyl,
15 methoxybenzyl, tetrahydrofuranylmethyl and (ethoxycarbonyl)ethyl.

36. A compound according to claim 33 wherein R₂ is chosen from hydrogen, lower alkyl and substituted lower alkyl, and R₂' is hydrogen.

37. A compound according to claim 36 wherein R₂ is chosen from hydrogen, methyl, ethyl, propyl, methylthioethyl, aminobutyl, (CBZ)aminobutyl,
20 cyclohexylmethyl, benzyloxymethyl, methylsulfinylethyl, methylsulfinylmethyl, hydroxymethyl, benzyl and indolylmethyl.

38. A compound according to claim 33 wherein R₃ is chosen from C₁-C₁₃

alkyl; substituted lower alkyl; phenyl; naphthyl; phenyl substituted with one or more halo, lower alkyl, loweralkoxy, nitro, carboxy, methylenedioxy, or trifluoromethyl; biphenyl; benzyl; phenoxymethyl; halophenoxymethyl; phenylvinyl; heteroaryl; heteroaryl substituted with lower alkyl; and benzyloxymethyl.

- 5 39. A compound according to claim 38 wherein R_3 is chosen from ethyl, propyl, chloropropyl, butoxy, heptyl, butyl, octyl, tridecanyl, (ethoxycarbonyl)ethyl, dimethylaminoethyl, dimethylaminomethyl, phenyl, naphthyl, halophenyl, dihalophenyl, cyanophenyl, halo(trifluoromethyl)phenyl, chlorophenoxymethyl, methoxyphenyl, carboxyphenyl, ethylphenyl, tolyl, biphenyl, methylenedioxyphenyl, 10 methylsulfonylphenyl, methoxychlorophenyl, chloronaphthyl, methylhalophenyl, trifluoromethylphenyl, butylphenyl, pentylphenyl, methylnitrophenyl, phenoxymethyl, dimethoxyphenyl, phenylvinyl, nitrochlorophenyl, nitrophenyl, dinitrophenyl, bis(trifluoromethyl)phenyl, benzyloxymethyl, benzyl, furanyl, benzofuranyl, pyridinyl, indolyl, methylpyridinyl, quinolinyl, picolinyl, pyrazolyl, and imidazolyl.

- 15 40. A compound according to claim 33 wherein R_3 is R_{15} -NH- and R_{15} is chosen from lower alkyl; cyclohexyl; phenyl; and phenyl substituted with halo, lower alkyl, loweralkoxy, or lower alkylthio.

41. A compound according to claim 40 wherein R_{15} is chosen from isopropyl, butyl, cyclohexyl, phenyl, bromophenyl, dichlorophenyl, methoxyphenyl, 20 ethylphenyl, tolyl, trifluoromethylphenyl and methylthiophenyl.

42. A compound according to claim 33 wherein R_4 is chosen from lower alkyl, substituted lower alkyl, cyclohexyl; phenyl substituted with hydroxy, lower alkoxy or lower alkyl; benzyl; heteroarylmethyl; heteroarylethyl; heteroarylpropyl and R_{16} -alkylene-, wherein R_{16} is amino, lower alkylamino, di(lower alkyl)amino, lower 25 alkoxy, or N-heterocyclyl.

43. A compound according to claim 42 wherein R_4 is chosen from methyl, ethyl, propyl, butyl, cyclohexyl, carboxyethyl, carboxymethyl, methoxyethyl, hydroxyethyl, hydroxypropyl, dimethylaminoethyl, dimethylaminopropyl,

- diethylaminoethyl, diethylaminopropyl, aminopropyl, methylaminopropyl, , 2,2-dimethyl-3-(dimethylamino)propyl, 1-cyclohexyl-4-(diethylamino)butyl, aminoethyl, aminobutyl, aminopentyl, aminohexyl, aminoethoxyethyl, isopropylaminopropyl, diisopropylaminoethyl, 1-methyl-4-(diethylamino)butyl, (t-Boc)aminopropyl,
- 5 hydroxyphenyl, benzyl, methoxyphenyl, methylmethoxyphenyl, dimethylphenyl, tolyl, ethylphenyl, (oxopyrrolidinyl)propyl, (methoxycarbonyl)ethyl, benzylpiperidinyl, pyridinylethyl, pyridinylmethyl, morpholinylethyl morpholinylpropyl, piperidinyl, azetidinylmethyl, azetidinypropyl pyrrolidinylethyl, pyrrolidinylpropyl, piperidinylmethyl, piperidinylethyl, imidazolylpropyl, imidazolylethyl,
- 10 (ethylpyrrolidinyl)methyl, (methylpyrrolidinyl)ethyl, (methylpiperidinyl)propyl, (methylpiperazinyl)propyl, furanylmethyl and indolylethyl.

44. A compound according to claim 33 wherein

R_1 is chosen from lower alkyl, benzyl, substituted benzyl and substituted phenyl;

R_2 is chosen from hydrogen, alkyl, substituted lower alkyl and benzyl;

15 R_2' is hydrogen;

R_3 is chosen from substituted phenyl and naphthyl;

R_4 is chosen from substituted alkyl and R_{16} -alkylene-;

R_5 is hydrogen or halo

R_6 is hydrogen, methyl or halo;

20 R_7 is hydrogen, halo, methyl or trifluoromethyl;

R_8 is hydrogen or halo;

R_{16} is chosen from di(lower alkylamino), (lower alkyl)amino, amino N-heterocyclyl and substituted N-heterocyclyl.

45. A compound according to claim 31 wherein

25 R_1 is benzyl or halobenzyl;

R_2 is chosen from ethyl and propyl;

R_2' is hydrogen;

R_3 is substituted phenyl;

$R_{3'}$ is substituted phenyl;

30 $R_{3''}$ is substituted phenyl;

R_4 is $(CH_2)_m OH$ or $(CH_2)_p R_{16}$ wherein m is 2 or 3 and p is 1-3;

R_5 is hydrogen;


R₆ is hydrogen;

R₇ is halo;

R₈ is hydrogen;

R₁₆ is chosen from amino, propylamino, and azetidiny.

- 5 46. A compound according to claim 31 of formula:



47. A compound according to claim 46 wherein:

R₁ is chosen from hydrogen, lower alkyl, substituted lower alkyl, benzyl, substituted benzyl, phenyl, naphthyl and substituted phenyl;

- 10 R₂ is chosen from hydrogen, lower alkyl and substituted lower alkyl and R₂' is hydrogen;

R₃' is chosen from C₁-C₁₃ alkyl; phenyl; naphthyl; phenyl substituted with halo, lower alkyl, lower alkoxy, nitro, methylenedioxy, or trifluoromethyl; biphenyl, benzyl and heteroaryl; and

- 15 R₄ is chosen from lower alkyl, substituted lower alkyl, cyclohexyl; phenyl substituted with hydroxy, lower alkoxy or lower alkyl; benzyl; heteroarylmethyl; heteroarylethyl; heteroarylpropyl and R₁₆-alkylene, wherein

R₁₆ is amino, (lower alkyl)amino, di(lower alkyl)amino, lower alkoxy, or N-heterocyclyl.

- 20 48. A compound according to claim 47 wherein

R₁ is chosen from lower alkyl, benzyl, substituted benzyl and substituted phenyl;

R₂ is hydrogen or lower alkyl;

R₂' is hydrogen;

R₃ is chosen from substituted phenyl and naphthyl;


R_4 is R_{16} -alkylene-, hydroxy(lower alkyl) or carboxy (lower alkyl);

R_7 is hydrogen, fluoro, chloro or methyl;

R_5 , R_6 and R_8 are hydrogen;

R_{16} is chosen from di(lower alkyl)amino, (lower alkyl)amino, amino, pyrrolidinyl and
5 piperidinyl.

49. A compound according to claim 31 of formula:



50. A compound according to claim 49 wherein:

R_1 is chosen from hydrogen, lower alkyl, substituted lower alkyl, benzyl, substituted
10 benzyl, phenyl, naphthyl and substituted phenyl;

R_2 is chosen from hydrogen, lower alkyl and substituted lower alkyl and R_2' is
hydrogen; and

R_4 is chosen from lower alkyl, cyclohexyl; phenyl substituted with hydroxy, lower
alkoxy or lower alkyl; benzyl; heteroarylmethyl; heteroarylethyl;
15 heteroarylpropyl and R_{16} -alkylene, wherein R_{16} is di(lower alkyl)amino,
(lower alkyl)amino, amino, lower alkoxy, or N-heterocyclyl.

51. A compound according to claim 49 wherein

R_1 is chosen from lower alkyl, benzyl, substituted benzyl and substituted phenyl;

R_2 is hydrogen or lower alkyl;

20 R_2' is hydrogen;


R_4 is R_{16} -alkylene-;

R_7 is hydrogen, fluoro, chloro or methyl;

R_5 , R_6 and R_8 are hydrogen;

R_{16} is chosen from di(lower alkylamino), (lower alkyl)amino, amino, pyrrolidinyl, piperidinyl, imidazolyl and morpholinyl.

52. A compound according to claim 31 of formula:



5 53. A compound according to claim 52 wherein:

R_1 is chosen from hydrogen, lower alkyl, substituted lower alkyl, benzyl, substituted benzyl, phenyl, naphthyl and substituted phenyl;

R_2 is chosen from hydrogen, lower alkyl and substituted lower alkyl and R_2' is hydrogen;

10 $R_{3''}$ is chosen from C_1 - C_{13} alkyl; substituted lower alkyl; phenyl; naphthyl; phenyl substituted with halo, lower alkyl, lower alkoxy, nitro, methylenedioxy, or trifluoromethyl; biphenyl, benzyl and heterocyclyl; and

R_4 is chosen from lower alkyl, substituted lower alkyl, cyclohexyl; phenyl substituted with hydroxy, lower alkoxy or lower alkyl; benzyl; substituted benzyl, heterocyclyl, heteroarylmethyl; heteroarylethyl; heteroarylpropyl and R_{16} -alkylene, wherein

R_{16} is di(lower alkyl)amino, (lower alkyl)amino, amino, lower alkoxy, or N-heterocyclyl.

54. A compound according to claim 53 wherein

20 R_1 is chosen from lower alkyl, benzyl, substituted benzyl and substituted phenyl;

R_2 is hydrogen or lower alkyl;

R_2' is hydrogen;

$R_{3''}$ is chosen from substituted phenyl, heterocyclyl and naphthyl;

R₄ is chosen from substituted benzyl, heterocyclyl substituted lower alkyl and R₁₆-alkylene-;

R₆ and R₇ are chosen from hydrogen and halo;

R₅ and R₈ are hydrogen;

- 5 R₁₆ is chosen from di(lower alkylamino), (lower alkyl)amino, amino, pyrrolidino, piperidino, imidazolyl and morpholino.

55. A compound according to claim 54 wherein

R₁ is benzyl;

R₂ is ethyl;

- 10 R₂' is hydrogen;

R₃ is chosen from halophenyl, polyhalophenyl, tolyl, dimethylphenyl, methoxyphenyl, dimethoxyphenyl, cyanophenyl, trifluoromethylphenyl, trifluoromethoxyphenyl, bis(trifluoromethyl)phenyl, carboxyphenyl, t-butylphenyl, methoxycarbonylphenyl, piperidinyl and naphthyl;

- 15 R₄ is chosen from substituted benzyl, piperidinyl, hydroxy(lower alkyl) and R₁₆-alkylene-;

R₆ and R₇ are chosen from hydrogen and halo;

R₅ and R₈ are hydrogen;




R₁₆ is chosen from dimethylamino, amino, pyrrolidinyl and piperidinyl.

- 20 56. A compound according to any of claims 32 to 45, 47, 48, 50, 51 and 53 to 55 wherein the stereogenic center to which R₂ and R₂' are attached is of the R configuration.


- 25 57. A compound according to claim 31 wherein said compound is of a formula as defined in Figure 3.

58. A method of screening for KSP kinesin modulators comprising:

- (a) combining a kinesin, a candidate bioactive agent and a compound chosen from the group consisting of:



and



wherein:

- 5 R_1 is chosen from hydrogen, alkyl, aryl, alkylaryl, heteroaryl, alkylheteroaryl, substituted alkyl, substituted aryl, substituted alkylaryl, substituted heteroaryl, and substituted alkylheteroaryl;
- R_2 and R_2' are independently chosen from hydrogen, alkyl, oxaalkyl, aryl, alkylaryl, heteroaryl, alkylheteroaryl, substituted alkyl, substituted aryl, substituted alkylaryl, substituted heteroaryl, and substituted alkylheteroaryl; or R_2 and R_2' taken together form a 3- to 7-membered ring;
- R_3 is chosen from hydrogen, alkyl, aryl, alkylaryl, heteroaryl, alkylheteroaryl, substituted alkyl, substituted aryl, substituted alkylaryl, substituted heteroaryl, substituted alkylheteroaryl, oxaalkyl, oxaalkylaryl, substituted oxaalkylaryl, $R_{15}O-$ and $R_{15}-NH-$;
- 15 $R_{3'}$ is chosen from hydrogen, alkyl, aryl, alkylaryl, heteroaryl, alkylheteroaryl, substituted alkyl, substituted aryl, substituted alkylaryl, substituted heteroaryl, substituted alkylheteroaryl and $R_{15}-NH-$;
- $R_{3''}$ is chosen from alkyl, aryl, alkylaryl, heteroaryl, alkylheteroaryl, substituted alkyl, substituted aryl, substituted alkylaryl, substituted heteroaryl, and substituted
- 20

alkylheteroaryl;

R₄ is chosen from hydrogen, alkyl, aryl, alkylaryl, heteroaryl, alkylheteroaryl, substituted alkyl, substituted aryl, substituted alkylaryl, substituted heteroaryl, substituted alkylheteroaryl, and R₁₆-alkylene-;

- 5 R₅, R₆, R₇ and R₈ are independently chosen from hydrogen, alkyl, alkoxy, halogen, fluoroalkyl, nitro, dialkylamino, alkylsulfonyl, alkylsulfonamido, sulfonamidoalkyl, sulfonamidoaryl, alkylthio, carboxyalkyl, carboxamido, aminocarbonyl, aryl and heteroaryl;



10 R₁₅ is chosen from hydrogen, alkyl, aryl, alkylaryl, heteroaryl, alkylheteroaryl, substituted alkyl, substituted aryl, substituted alkylaryl, substituted heteroaryl, and substituted alkylheteroaryl;

R₁₆ is chosen from alkoxy, amino, alkylamino, dialkylamino, N-heterocyclyl and substituted N-heterocyclyl; and


- (b) determining the effect of said candidate bioactive agent on the
15 activity of said kinesin.

59. A method of screening for compounds that bind to KSP kinesin comprising:

- (a) combining a kinesin, a candidate bioactive agent and a labeled compound chosen from the group consisting of:



20



wherein:

R_1 is chosen from hydrogen, alkyl, aryl, alkylaryl, heteroaryl, alkylheteroaryl, substituted alkyl, substituted aryl, substituted alkylaryl, substituted heteroaryl, and substituted alkylheteroaryl;

R_2 and R_2' are independently chosen from hydrogen, alkyl, oxaalkyl, aryl, alkylaryl, heteroaryl, alkylheteroaryl, substituted alkyl, substituted aryl, substituted alkylaryl, substituted heteroaryl, and substituted alkylheteroaryl; or R_2 and R_2' taken together form a 3- to 7-membered ring;

R_3 is chosen from hydrogen, alkyl, aryl, alkylaryl, heteroaryl, alkylheteroaryl, substituted alkyl, substituted aryl, substituted alkylaryl, substituted heteroaryl, substituted alkylheteroaryl, oxaalkyl, oxaalkylaryl, substituted oxaalkylaryl, $R_{15}O-$ and $R_{15}NH-$;

R_3' is chosen from hydrogen, alkyl, aryl, alkylaryl, heteroaryl, alkylheteroaryl, substituted alkyl, substituted aryl, substituted alkylaryl, substituted heteroaryl, substituted alkylheteroaryl and $R_{15}NH-$;

R_3'' is chosen from alkyl, aryl, alkylaryl, heteroaryl, alkylheteroaryl, substituted alkyl, substituted aryl, substituted alkylaryl, substituted heteroaryl, and substituted alkylheteroaryl;

R_4 is chosen from hydrogen, alkyl, aryl, alkylaryl, heteroaryl, alkylheteroaryl, substituted alkyl, substituted aryl, substituted alkylaryl, substituted heteroaryl, substituted alkylheteroaryl, and R_{16} -alkylene-;

R_5 , R_6 , R_7 and R_8 are independently chosen from hydrogen, alkyl, alkoxy, halogen, fluoroalkyl, nitro, dialkylamino, alkylsulfonyl, alkylsulfonamido, sulfonamidoalkyl, sulfonamidoaryl, alkylthio, carboxyalkyl, carboxamido,

aminocarbonyl, aryl and heteroaryl;

R₁₅ is chosen from hydrogen, alkyl, aryl, alkylaryl, heteroaryl, alkylheteroaryl, substituted alkyl, substituted aryl, substituted alkylaryl, substituted heteroaryl, and substituted alkylheteroaryl;

- 5 R₁₆ is chosen from alkoxy, amino, alkylamino, dialkylamino, N-heterocyclyl and substituted N-heterocyclyl; and

(b) determining the binding of said candidate bioactive agent to said kinesin.

Figure 1

5




Figure 2





Figure 3



R ₁	R ₂	R ₃	R ₄	R ₅	R ₆	R ₇	R ₈	Activity
						Cl-X ₇		%I > 35% @ 40 uM
								%I > 35% @ 40 uM
								%I > 35% @ 40 uM

Figure 3 (continued)

R ₁	R ₂	R ₃	R ₄	R ₅	R ₆	R ₇	R ₈	Activity
								%I > 35% @ 40 uM
								%I > 35% @ 40 uM
								%I > 35% @ 40 uM
								%I > 35% @ 40 uM
								%I > 35% @ 40 uM

Figure 3 (continued)

R ₁	R ₂	R ₃	R ₄	R ₅	R ₆	R ₇	R ₈	Activity
								%I > 35% @ 40 uM
								%I > 35% @ 40 uM
								%I > 35% @ 40 uM
								%I > 35% @ 40 uM
								%I > 35% @ 40 uM

Figure 3 (continued)
























R ₁	R ₂	R ₃	R ₄	R ₅	R ₆	R ₇	R ₈	Activity
								%I > 35% @ 40 uM
								%I > 35% @ 40 uM
								%I > 35% @ 40 uM
								%I > 35% @ 40 uM
								%I > 35% @ 40 uM

Figure 3 (continued)

R ₁	R ₂	R ₃	R ₄	R ₅	R ₆	R ₇	R ₈	Activity
								%I > 35% @ 40 uM
								%I > 35% @ 40 uM
								%I > 35% @ 40 uM
								%I > 35% @ 40 uM
								%I > 35% @ 40 uM

Figure 3 (continued)

















R ₁	R ₂	R ₃	R ₄	R ₅	R ₆	R ₇	R ₈	Activity
								%I > 35% @ 40 uM
								%I > 35% @ 40 uM
								%I > 35% @ 40 uM
								%I > 35% @ 40 uM

Figure 3 (continued)

















R ₁	R ₂	R ₃	R ₄	R ₅	R ₆	R ₇	R ₈	Activity
								%I > 35% @ 40 uM
								%I > 35% @ 40 uM
								%I > 35% @ 40 uM
								%I > 35% @ 40 uM

Figure 3 (continued)

R ₁	R ₂	R ₃	R ₄	R ₅	R ₆	R ₇	R ₈	Activity
								%I > 35% @ 40 uM
								%I > 35% @ 40 uM
								%I > 35% @ 40 uM
								%I > 35% @ 40 uM
								%I > 35% @ 40 uM

Figure 3 (continued)

R ₁	R ₂	R ₃	R ₄	R ₅	R ₆	R ₇	R ₈	Activity
	H ₃ C-X ₂							%I > 35% @ 40 uM
	H ₃ C-X ₂							%I > 35% @ 40 uM
	H ₃ C-X ₂							%I > 35% @ 40 uM
	H ₃ C-X ₂							%I > 35% @ 40 uM
	X ₂ -CH ₂ -CH ₃							%I > 35% @ 40 uM

Figure 3 (continued)

R ₁	R ₂	R ₃	R ₄	R ₅	R ₆	R ₇	R ₈	Activity
								%I > 35% @ 40 uM
								%I > 35% @ 40 uM
								%I > 35% @ 40 uM
								%I > 35% @ 40 uM
								%I > 35% @ 40 uM

Figure 3 (continued)

R ₁	R ₂	R ₃	R ₄	R ₅	R ₆	R ₇	R ₈	Activity
								%I > 35% @ 40 uM
								%I > 35% @ 40 uM
								%I > 35% @ 40 uM
								%I > 35% @ 40 uM
								%I > 35% @ 40 uM

Figure 3 (continued)





















R ₁	R ₂	R ₃	R ₄	R ₅	R ₆	R ₇	R ₈	Activity
								%I > 35% @ 40 uM
								%I > 35% @ 40 uM
								%I > 35% @ 40 uM
								%I > 35% @ 40 uM

Figure 3 (continued)



















R ₁	R ₂	R ₃	R ₄	R ₅	R ₆	R ₇	R ₈	Activity
								%I > 35% @ 40 uM
								%I > 35% @ 40 uM
								%I > 35% @ 40 uM
								%I > 35% @ 40 uM

Figure 3 (continued)

R ₁	R ₂	R ₃	R ₄	R ₅	R ₆	R ₇	R ₈	Activity
								%I > 35% @ 40 uM
								%I > 35% @ 40 uM
								%I > 35% @ 40 uM
								%I > 35% @ 40 uM
								%I > 35% @ 40 uM

Figure 3 (continued)













R ₁	R ₂	R ₃	R ₄	R ₅	R ₆	R ₇	R ₈	Activity
H ₃ C-X ₁	H ₃ C-X ₂					X ₇ -Cl		%I > 35% @ 40 uM
H ₃ C-X ₁	H ₃ C-X ₂					X ₇ -Cl		%I > 35% @ 40 uM
								%I > 35% @ 40 uM
								%I > 35% @ 40 uM

Figure 3 (continued)

















R ₁	R ₂	R ₃	R ₄	R ₅	R ₆	R ₇	R ₈	Activity
								%I > 35% @ 40 uM
								%I > 35% @ 40 uM
								%I > 35% @ 40 uM
								%I > 35% @ 40 uM

Figure 3 (continued)

R ₁	R ₂	R ₃	R ₄	R ₅	R ₆	R ₇	R ₈	Activity
								%I > 35% @ 40 uM
								%I > 35% @ 40 uM
								%I > 35% @ 40 uM
								%I > 35% @ 40 uM

Figure 3 (continued)

R ₁	R ₂	R ₃	R ₄	R ₅	R ₆	R ₇	R ₈	Activity
								%I > 35% @ 40 uM
								%I > 35% @ 40 uM
								%I > 35% @ 40 uM
								%I > 35% @ 40 uM
								%I > 35% @ 40 uM

Figure 3 (continued)

R ₁	R ₂	R ₃	R ₄	R ₅	R ₆	R ₇	R ₈	Activity
								%I > 35% @ 40 uM
								%I > 35% @ 40 uM
								%I > 35% @ 40 uM
								%I > 35% @ 40 uM
								%I > 35% @ 40 uM

Figure 3 (continued)

















R ₁	R ₂	R ₃	R ₄	R ₅	R ₆	R ₇	R ₈	Activity
								%I > 35% @ 40 uM
								%I > 35% @ 40 uM
								%I > 35% @ 40 uM
								%I > 35% @ 40 uM

Figure 3 (continued)

















R ₁	R ₂	R ₃	R ₄	R ₅	R ₆	R ₇	R ₈	Activity
								%I > 35% @ 40 uM
								%I > 35% @ 40 uM
								%I > 35% @ 40 uM
								%I > 35% @ 40 uM

Figure 3 (continued)













R ₁	R ₂	R ₃	R ₄	R ₅	R ₆	R ₇	R ₈	Activity
	$\text{H}_3\text{C}-\text{X}_2$					X_7-Cl		%I > 35% @ 40 uM
	X_2-CH_3					$\text{Cl}-\text{X}_7$		%I > 35% @ 40 uM
	$\text{H}_3\text{C}-\text{X}_2$					X_7-Cl		%I > 35% @ 40 uM
	$\text{H}_3\text{C}-\text{X}_2$							%I > 35% @ 40 uM

Figure 3 (continued)

















R ₁	R ₂	R ₃	R ₄	R ₅	R ₆	R ₇	R ₈	Activity
								%I > 35% @ 40 uM
								%I > 35% @ 40 uM
								%I > 35% @ 40 uM
								%I > 35% @ 40 uM

Figure 3 (continued)

R ₁	R ₂	R ₃	R ₄	R ₅	R ₆	R ₇	R ₈	Activity
								%I > 35% @ 40 uM
								%I > 35% @ 40 uM
								%I > 35% @ 40 uM
								%I > 35% @ 40 uM
								%I > 35% @ 40 uM

Figure 3 (continued)

















R ₁	R ₂	R ₃	R ₄	R ₅	R ₆	R ₇	R ₈	Activity
								%I > 35% @ 40 uM
								%I > 35% @ 40 uM
								%I > 35% @ 40 uM
								%I > 35% @ 40 uM

Figure 3 (continued)

















R ₁	R ₂	R ₃	R ₄	R ₅	R ₆	R ₇	R ₈	Activity
								%I > 35% @ 40 uM
								%I > 35% @ 40 uM
								%I > 35% @ 40 uM
								%I > 35% @ 40 uM

Figure 3 (continued)

















R ₁	R ₂	R ₃	R ₄	R ₅	R ₆	R ₇	R ₈	Activity
								%I > 35% @ 40 uM
								%I > 35% @ 40 uM
								%I > 35% @ 40 uM
								%I > 35% @ 40 uM

Figure 3 (continued)









R ₁	R ₂	R ₃	R ₄	R ₅	R ₆	R ₇	R ₈	Activity
								%I > 35% @ 40 uM
								%I > 35% @ 40 uM

Figure 3 (continued)

















								
R ₁	R ₂	R ₃	R ₄	R ₅	R ₆	R ₇	R ₈	Activity
								IC50 < 100 nM
								IC50 < 100 nM
								IC50 < 100 nM

Figure 3 (continued)


























R ₁	R ₂	R ₃	R ₄	R ₅	R ₆	R ₇	R ₈	Activity
								IC50 < 100 nM
								IC50 < 100 nM
								IC50 < 100 nM
								IC50 < 100 nM
								IC50 < 100 nM

Figure 3 (continued)


























R ₁	R ₂	R ₃	R ₄	R ₅	R ₆	R ₇	R ₈	Activity
								IC50 < 100 nM
								IC50 < 100 nM
								IC50 < 100 nM
								IC50 < 100 nM
								IC50 < 100 nM

Figure 3 (continued)


























R ₁	R ₂	R ₃	R ₄	R ₅	R ₆	R ₇	R ₈	Activity
								IC50 < 100 nM
								IC50 < 100 nM
								IC50 < 100 nM
								IC50 < 100 nM
								IC50 < 100 nM

Figure 3 (continued)

R ₁	R ₂	R ₃	R ₄	R ₅	R ₆	R ₇	R ₈	Activity
								IC50 < 100 nM
								IC50 < 100 nM
								IC50 < 100 nM
								IC50 < 100 nM
								IC50 < 100 nM

Figure 3 (continued)

R ₁	R ₂	R ₃	R ₄	R ₅	R ₆	R ₇	R ₈	Activity
								IC50 < 100 nM
								IC50 < 100 nM
								IC50 < 100 nM
								IC50 < 100 nM
								IC50 < 100 nM

Figure 3 (continued)


























R ₁	R ₂	R ₃	R ₄	R ₅	R ₆	R ₇	R ₈	Activity
								IC50 < 100 nM
								IC50 < 100 nM
								IC50 < 100 nM
								IC50 < 100 nM
								IC50 < 100 nM

Figure 3 (continued)

R ₁	R ₂	R ₃	R ₄	R ₅	R ₆	R ₇	R ₈	Activity
								IC50 < 100 nM
								IC50 < 100 nM
								IC50 < 100 nM
								IC50 < 100 nM
								IC50 < 100 nM

Figure 3 (continued)


























R ₁	R ₂	R ₃	R ₄	R ₅	R ₆	R ₇	R ₈	Activity
								IC50 < 100 nM
								IC50 < 100 nM
								IC50 < 100 nM
								IC50 < 100 nM
								IC50 < 100 nM

Figure 3 (continued)

R ₁	R ₂	R ₃	R ₄	R ₅	R ₆	R ₇	R ₈	Activity
								IC50 < 100 nM
								IC50 < 100 nM
								IC50 < 100 nM
								IC50 < 100 nM
								IC50 < 100 nM

Figure 3 (continued)

R ₁	R ₂	R ₃	R ₄	R ₅	R ₆	R ₇	R ₈	Activity
								IC50 < 100 nM
								IC50 < 100 nM
								IC50 < 100 nM
								IC50 < 100 nM
								IC50 < 100 nM

Figure 3 (continued)

R ₁	R ₂	R ₃	R ₄	R ₅	R ₆	R ₇	R ₈	Activity
								IC50 < 100 nM
								IC50 < 100 nM
								IC50 < 100 nM
								IC50 < 100 nM
				40				IC50 < 100 nM

Figure 3 (continued)


























R ₁	R ₂	R ₃	R ₄	R ₅	R ₆	R ₇	R ₈	Activity
								IC50 < 100 nM
								IC50 < 100 nM
								IC50 < 100 nM
								IC50 < 100 nM
								IC50 < 100 nM

Figure 3 (continued)

R ₁	R ₂	R ₃	R ₄	R ₅	R ₆	R ₇	R ₈	Activity
								IC50 < 100 nM
								IC50 < 100 nM
								IC50 < 100 nM
								IC50 < 100 nM
								IC50 < 100 nM

Figure 3 (continued)

R ₁	R ₂	R ₃	R ₄	R ₅	R ₆	R ₇	R ₈	Activity
								IC50 < 100 nM
								IC50 < 100 nM
								IC50 < 100 nM
								IC50 < 100 nM
								IC50 < 100 nM

Figure 3 (continued)

R ₁	R ₂	R ₃	R ₄	R ₅	R ₆	R ₇	R ₈	Activity
								IC50 < 100 nM
								IC50 < 100 nM
								IC50 = 100 nM-1 uM
								IC50 = 100 nM-1 uM
								IC50 = 100 nM-1 uM

Figure 3 (continued)

R ₁	R ₂	R ₃	R ₄	R ₅	R ₆	R ₇	R ₈	Activity
								IC50 = 100 nM-1 uM
								IC50 = 100 nM-1 uM
								IC50 = 100 nM-1 uM
								IC50 = 100 nM-1 uM
								IC50 = 100 nM-1 uM

Figure 3 (continued)

R ₁	R ₂	R ₃	R ₄	R ₅	R ₆	R ₇	R ₈	Activity
								IC50 = 100 nM-1 uM
								IC50 = 100 nM-1 uM
								IC50 = 100 nM-1 uM
								IC50 = 100 nM-1 uM
								IC50 = 100 nM-1 uM

Figure 3 (continued)

R ₁	R ₂	R ₃	R ₄	R ₅	R ₆	R ₇	R ₈	Activity
								IC50 = 100 nM-1 uM
								IC50 = 100 nM-1 uM
								IC50 = 100 nM-1 uM
								IC50 = 100 nM-1 uM
								IC50 = 100 nM-1 uM

Figure 3 (continued)


























R ₁	R ₂	R ₃	R ₄	R ₅	R ₆	R ₇	R ₈	Activity
								IC50 = 100 nM-1 uM
								IC50 = 100 nM-1 uM
								IC50 = 100 nM-1 uM
								IC50 = 100 nM-1 uM
								IC50 = 100 nM-1 uM

Figure 3 (continued)

R ₁	R ₂	R ₃	R ₄	R ₅	R ₆	R ₇	R ₈	Activity
								IC50 = 100 nM-1 uM
								IC50 = 100 nM-1 uM
								IC50 = 100 nM-1 uM
								IC50 = 100 nM-1 uM
								IC50 = 100 nM-1 uM

Figure 3 (continued)

R ₁	R ₂	R ₃	R ₄	R ₅	R ₆	R ₇	R ₈	Activity
								IC50 = 100 nM-1 uM
								IC50 = 100 nM-1 uM
								IC50 = 100 nM-1 uM
								IC50 = 100 nM-1 uM
								IC50 = 100 nM-1 uM

Figure 3 (continued)

R ₁	R ₂	R ₃	R ₄	R ₅	R ₆	R ₇	R ₈	Activity
								IC50 = 100 nM-1 uM
								IC50 = 100 nM-1 uM
								IC50 = 100 nM-1 uM
								IC50 = 100 nM-1 uM
								IC50 = 100 nM-1 uM

Figure 3 (continued)

R ₁	R ₂	R ₃	R ₄	R ₅	R ₆	R ₇	R ₈	Activity
								IC50 = 100 nM-1 uM
								IC50 = 100 nM-1 uM
								IC50 = 100 nM-1 uM
								IC50 = 100 nM-1 uM

Figure 3 (continued)

R ₁	R ₂	R ₃	R ₄	R ₅	R ₆	R ₇	R ₈	Activity
								IC50 = 100 nM-1 uM
								IC50 = 100 nM-1 uM
								IC50 = 100 nM-1 uM
								IC50 = 100 nM-1 uM
								IC50 = 100 nM-1 uM

Figure 3 (continued)

























R ₁	R ₂	R ₃	R ₄	R ₅	R ₆	R ₇	R ₈	Activity
								IC50 = 100 nM-1 uM
								IC50 = 100 nM-1 uM
								IC50 = 100 nM-1 uM
								IC50 = 100 nM-1 uM
								IC50 = 100 nM-1 uM

Figure 3 (continued)

R ₁	R ₂	R ₃	R ₄	R ₅	R ₆	R ₇	R ₈	Activity
								IC50 = 100 nM-1 uM
								IC50 = 100 nM-1 uM
								IC50 = 100 nM-1 uM
								IC50 = 100 nM-1 uM
								IC50 = 100 nM-1 uM

Figure 3 (continued)

R ₁	R ₂	R ₃	R ₄	R ₅	R ₆	R ₇	R ₈	Activity
								IC50 = 100 nM-1 uM
								IC50 = 100 nM-1 uM
								IC50 = 100 nM-1 uM
								IC50 = 100 nM-1 uM
								IC50 = 1 uM-10 uM

Figure 3 (continued)

R ₁	R ₂	R ₃	R ₄	R ₅	R ₆	R ₇	R ₈	Activity
								IC50 = 1 uM-10 uM
								IC50 = 1 uM-10 uM
								IC50 = 1 uM-10 uM
								IC50 = 1 uM-10 uM
								IC50 = 1 uM-10 uM

Figure 3 (continued)





















R ₁	R ₂	R ₃	R ₄	R ₅	R ₆	R ₇	R ₈	Activity
								IC50 = 1 uM-10 uM
								IC50 = 1 uM-10 uM
								IC50 = 1 uM-10 uM
								IC50 = 1 uM-10 uM

Figure 3 (continued)




















R ₁	R ₂	R ₃	R ₄	R ₅	R ₆	R ₇	R ₈	Activity
								IC50 = 1 uM-10 uM
								IC50 = 1 uM-10 uM
								IC50 = 1 uM-10 uM
								IC50 = 1 uM-10 uM

Figure 3 (continued)


















R ₁	R ₂	R ₃	R ₄	R ₅	R ₆	R ₇	R ₈	Activity
								IC50 = 1 uM-10 uM
								IC50 = 1 uM-10 uM
								IC50 = 1 uM-10 uM
								IC50 = 1 uM-10 uM

Figure 3 (continued)

R ₁	R ₂	R ₃	R ₄	R ₅	R ₆	R ₇	R ₈	Activity
								IC50 = 1 uM-10 uM
								IC50 = 1 uM-10 uM
								IC50 = 1 uM-10 uM
								IC50 = 1 uM-10 uM
								IC50 = 1 uM-10 uM

Figure 3 (continued)






















R ₁	R ₂	R ₃	R ₄	R ₅	R ₆	R ₇	R ₈	Activity
								IC50 = 1 uM-10 uM
								IC50 = 1 uM-10 uM
								IC50 = 1 uM-10 uM
								IC50 = 1 uM-10 uM
								IC50 = 1 uM-10 uM

Figure 3 (continued)

























R ₁	R ₂	R ₃	R ₄	R ₅	R ₆	R ₇	R ₈	Activity
								IC50 = 1 uM-10 uM
								IC50 = 1 uM-10 uM
								IC50 = 1 uM-10 uM
								IC50 = 1 uM-10 uM
								IC50 = 1 uM-10 uM

Figure 3 (continued)






















R ₁	R ₂	R ₃	R ₄	R ₅	R ₆	R ₇	R ₈	Activity
								IC50 = 1 uM-10 uM
								IC50 = 1 uM-10 uM
								IC50 = 1 uM-10 uM
								IC50 = 1 uM-10 uM
								IC50 = 1 uM-10 uM

Figure 3 (continued)

R ₁	R ₂	R ₃	R ₄	R ₅	R ₆	R ₇	R ₈	Activity
								IC50 = 1 uM-10 uM
								IC50 = 1 uM-10 uM
								IC50 = 1 uM-10 uM
								IC50 = 1 uM-10 uM
								IC50 = 1 uM-10 uM

Figure 3 (continued)

R ₁	R ₂	R ₃	R ₄	R ₅	R ₆	R ₇	R ₈	Activity
								IC50 = 1 uM-10 uM
								IC50 = 1 uM-10 uM
								IC50 = 1 uM-10 uM
								IC50 = 1 uM-10 uM
								IC50 = 1 uM-10 uM

Figure 3 (continued)

R ₁	R ₂	R ₃	R ₄	R ₅	R ₆	R ₇	R ₈	Activity
								IC50 = 1 uM-10 uM
								IC50 = 1 uM-10 uM
								IC50 = 1 uM-10 uM
								IC50 = 10 uM-50 uM
								IC50 = 10 uM-50 uM

Figure 3 (continued)























R ₁	R ₂	R ₃	R ₄	R ₅	R ₆	R ₇	R ₈	Activity
								IC50 = 10 uM-50 uM
								IC50 = 10 uM-50 uM
								IC50 = 10 uM-50 uM
								IC50 = 10 uM-50 uM
								IC50 = 10 uM-50 uM

Figure 3 (continued)























R ₁	R ₂	R ₃	R ₄	R ₅	R ₆	R ₇	R ₈	Activity
								IC50 = 10 uM-50 uM
								IC50 = 10 uM-50 uM
								IC50 = 10 uM-50 uM
								IC50 = 10 uM-50 uM
								IC50 = 10 uM-50 uM

Figure 3 (continued)























R ₁	R ₂	R ₃	R ₄	R ₅	R ₆	R ₇	R ₈	Activity
								IC50 = 10 uM-50 uM
								IC50 = 10 uM-50 uM
								IC50 = 10 uM-50 uM
								IC50 = 10 uM-50 uM
								IC50 = 10 uM-50 uM

Figure 3 (continued)
























R ₁	R ₂	R ₃	R ₄	R ₅	R ₆	R ₇	R ₈	Activity
								IC50 = 10 uM-50 uM
								IC50 = 10 uM-50 uM
								IC50 = 10 uM-50 uM
								IC50 = 10 uM-50 uM
								IC50 = 10 uM-50 uM

Figure 3 (continued)

R ₁	R ₂	R ₃	R ₄	R ₅	R ₆	R ₇	R ₈	Activity
								IC50 = 10 uM-50 uM
								IC50 = 10 uM-50 uM
								IC50 = 10 uM-50 uM
								IC50 = 10 uM-50 uM
								IC50 = 10 uM-50 uM

Figure 3 (continued)























R ₁	R ₂	R ₃	R ₄	R ₅	R ₆	R ₇	R ₈	Activity
								IC50 = 10 uM-50 uM
								IC50 = 10 uM-50 uM
								IC50 = 10 uM-50 uM
								IC50 = 10 uM-50 uM
								IC50 = 10 uM-50 uM

Figure 3 (continued)


























R ₁	R ₂	R ₃	R ₄	R ₅	R ₆	R ₇	R ₈	Activity
								IC50 = 10 uM-50 uM
								IC50 = 10 uM-50 uM
								IC50 = 10 uM-50 uM
								IC50 = 10 uM-50 uM
								IC50 = 10 uM-50 uM

Figure 3 (continued)



R ₁	R ₂	R ₃	R ₄	R ₅	R ₆	R ₇	R ₈	Activity
								IC50 = 10 uM-50 uM

Figure 3 (continued)

















								
R ₁	R ₂	R ₃ ''	R ₄	R ₅	R ₆	R ₇	R ₈	Activity
								Ki < 100 nM
								Ki < 100 nM
								Ki < 100 nM

Figure 3 (continued)

R ₁	R ₂	R ₃ ^{''}	R ₄	R ₅	R ₆	R ₇	R ₈	Activity
								Ki < 100 nM
								Ki < 100 nM
								Ki < 100 nM
								Ki < 100 nM
								Ki < 100 nM

Figure 3 (continued)


























R ₁	R ₂	R ₃ ^{''}	R ₄	R ₅	R ₆	R ₇	R ₈	Activity
								Ki < 100 nM
								Ki < 100 nM
								Ki < 100 nM
								Ki < 100 nM
								Ki < 100 nM

Figure 3 (continued)


























R ₁	R ₂	R ₃ "	R ₄	R ₅	R ₆	R ₇	R ₈	Activity
								Ki < 100 nM
								Ki < 100 nM
								Ki < 100 nM
								Ki < 100 nM
								Ki < 100 nM

Figure 3 (continued)

R ₁	R ₂	R ₃ "	R ₄	R ₅	R ₆	R ₇	R ₈	Activity
								Ki < 100 nM
								Ki < 100 nM
								Ki < 100 nM
								Ki < 100 nM
								Ki < 100 nM

Figure 3 (continued)

R ₁	R ₂	R ₃ "	R ₄	R ₅	R ₆	R ₇	R ₈	Activity
								Ki < 100 nM
								Ki < 100 nM
								Ki < 100 nM
								Ki < 100 nM
								Ki < 100 nM

Figure 3 (continued)


























R ₁	R ₂	R ₃ "	R ₄	R ₅	R ₆	R ₇	R ₈	Activity
								Ki < 100 nM
								Ki < 100 nM
								Ki < 100 nM
								Ki < 100 nM
								Ki < 100 nM

Figure 3 (continued)

R ₁	R ₂	R ₃ "	R ₄	R ₅	R ₆	R ₇	R ₈	Activity
								Ki < 100 nM
								Ki < 100 nM
								Ki < 100 nM
								Ki < 100 nM
								Ki < 100 nM

Figure 3 (continued)


























R ₁	R ₂	R ₃ ^{''}	R ₄	R ₅	R ₆	R ₇	R ₈	Activity
								Ki < 100 nM
								Ki < 100 nM
								Ki < 100 nM
								Ki < 100 nM
								Ki < 100 nM

Figure 3 (continued)

R ₁	R ₂	R ₃ "	R ₄	R ₅	R ₆	R ₇	R ₈	Activity
								Ki < 100 nM
								Ki < 100 nM
								Ki < 100 nM
								Ki = 100 nM-1 uM
								Ki = 100 nM-1 uM

Figure 3 (continued)

R ₁	R ₂	R ₃ "	R ₄	R ₅	R ₆	R ₇	R ₈	Activity
								Ki = 100 nM-1 uM
								Ki = 100 nM-1 uM
								Ki = 100 nM-1 uM
								Ki = 1 uM-10 uM
								Ki = 1 uM-10 uM

Figure 4: Asymmetric Synthesis






Figure 5

Figure 5a: Sulfonamide Synthesis





5

Figure 5b: Carbamate Synthesis



10

Figure 5c: Urea Synthesis

15 Figure 5d: Primary and Secondary Amine R_4 Synthesis

INTERNATIONAL SEARCH REPORT

Intern Application No
PCT/US 00/29585

A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 C07D239/91 A61K31/517 A61P37/02

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 C07D A61K A61P

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

PAJ, CHEM ABS Data

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	A.K. DEBNATH: "STRUCTURE BASED IDENTIFICATION OF SMALL MOLECULE ANTIVIRAL COMPOUNDS" JOURNAL OF MEDICINAL CHEMISTRY., vol. 42, no. 17, 26 August 1999 (1999-08-26), pages 3203-9, XP002159846 AMERICAN CHEMICAL SOCIETY. WASHINGTON., US ISSN: 0022-2623 page 3203 -page 3206 --- -/--	1, 31, 49, 58

☒ Further documents are listed in the continuation of box C.☒ Patent family members are listed in annex.

* Special categories of cited documents:

A document defining the general state of the art which is not considered to be of particular relevance

E earlier document but published on or after the international filing date

L document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

O document referring to an oral disclosure, use, exhibition or other means

P document published prior to the international filing date but later than the priority date claimed

T later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

X document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

Y document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

& document member of the same patent family

Date of the actual completion of the international search

8 February 2001

Date of mailing of the international search report

22/02/2001

Name and mailing address of the ISA

European Patent Office, P.B. 5618 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Francois, J

INTERNATIONAL SEARCH REPORT

Intern: Application No

PCT/US 00/29585

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	CHEMICAL ABSTRACTS, vol. 122, no. 26, 1995 Columbus, Ohio, US; abstract no. 327093n, BOCAKEI, ZAOLT: "TWO ANTITHROMBOTIC QUINAZOLONE DERIVATIVES." page 944; XP002159847 abstract & ACTA CRYSTALLOGR. SECT. C: CRYST. STRUCT. COMMUN., vol. C51, no. 4, 1995, pages 723-6, ---	1, 31, 49, 58
X	CHEMICAL ABSTRACTS, vol. 124, no. 13, 1996 Columbus, Ohio, US; abstract no. 176002v, SZABO, MONICA: "SYNTHESIS OF POTENTIAL CCK ANTAGONIST QUINAZOLONE." page 1287; column 1; XP002159848 abstract & ACTA PHARM. HUNG., vol. 65, no. 5, 1995, pages 175-81, HUNG	1, 31, 49, 58
A	EP 0 056 637 A (ISHIKAWA) 28 July 1982 (1982-07-28) cited in the application page 0; claims ---	1, 31, 58
A	EP 0 884 310 A (PFIZER) 16 December 1998 (1998-12-16) the whole document -----	1, 31, 49, 58

INTERNATIONAL SEARCH REPORT

Information on patent family members

Intern Application No
PCT/US 00/29585

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
EP 056637 A	28-07-1982	JP 57118570 A	23-07-1982
		JP 58023678 A	12-02-1983
		JP 58049370 A	23-03-1983
		AU 543928 B	09-05-1985
		AU 7917282 A	22-07-1982
		CA 1167442 A	15-05-1984
		DE 3264802 D	29-08-1985
		ES 508765 D	16-02-1983
		ES 8304097 A	16-05-1983
		US 4451467 A	29-05-1984
EP 884310 A	16-12-1998	BR 9801808 A	21-03-2000
		CA 2240138 A	09-12-1998
		JP 11012255 A	19-01-1999